

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
1 April 2004 (01.04.2004)

PCT

(10) International Publication Number
WO 2004/026120 A2

(51) International Patent Classification⁷: **A61B**
(21) International Application Number:
PCT/US2003/030213
(22) International Filing Date:
23 September 2003 (23.09.2003)
(25) Filing Language: English
(26) Publication Language: English
(30) Priority Data:
60/412,964 23 September 2002 (23.09.2002) US
(71) Applicant (for all designated States except US): **THE GENERAL HOSPITAL CORPORATION** [US/US]; 55 Fruit Street, Boston, MA 02114 (US).

(72) Inventors; and
(75) Inventors/Applicants (for US only): **SHELLEY, Carl**, Simon [GB/US]; 89 Myrtle Street #1, Boston, MA 02114 (US). **FAROKHZAD, Omid, C.** [US/US]; 84 Gainsborough Street #202W, Boston, MA 02115 (US).
(74) Agent: **LOCKHART, Helen, C.**, Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR DIAGNOSING AND TREATING TUMORS AND SUPPRESSING CD PROMOTERS



(57) Abstract: Methods of treating tumors, reducing white blood cell numbers, and inhibiting CD promoters are provided.

BEST AVAILABLE COPY

WO 2004/026120 A2

THIS PAGE BLANK (USPTO)

- 1 -

**METHODS FOR DIAGNOSING AND TREATING TUMORS AND SUPPRESSING
CD PROMOTERS**

FIELD OF THE INVENTION

5 The invention relates to methods of treating tumors and methods of reducing the number of white blood cells. The invention also relates to methods of diagnosing ovarian tumors and assessing the prognosis and the progression of disease in patients with ovarian tumors. The invention also relates to methods of inhibiting CD promoters.

10 **BACKGROUND OF THE INVENTION**

CD43 is a heavily glycosylated transmembrane molecule which plays a critical role in leukocyte activation and adhesion (Ostberg JR *et al.*, (1998) *Immunology Today*, 19:546-550; Remold-O'Donnell *et al.*, (1990) *Immunodeficiency Review*, 2:151-174; Rosenstein *et al.*, (1999) *Immunology Research*, 20:89-99). The importance of CD43 is
15 demonstrated by two immunodeficiency diseases, Wiskott-Aldrich syndrome (WAS) and the early stages of HIV infection (Parkman *et al.*, (1981) *Lancet*, ii:1387-1389; Remold-O'Donnell *et al.*, (1984) *Journal of Experimental Medicine*, 159:1705-1723; Ardman *et al.*, (1990) *Journal of Experimental Medicine*, 172:1151-1158; Lefebvre *et al.*, (1994) *Journal of Experimental Medicine*, 180:1609-1617; Giordanengo *et al.*, (1995)
20 *Blood*, 86:2302-2311; Gallego *et al.*, (2001) *AIDS*, 15:477-481). The etiology of both diseases involves the development of defects in CD43. Patients with WAS, which is an X-chromosome linked, recessive disorder, express defective CD43 on the surface of their T-lymphocytes. Affected males are subject to recurring opportunistic infections and do not respond to carbohydrate antigens reflecting defects in T-lymphocyte function. In addition,
25 patients suffer from eczema and thrombocytopenia with platelets of reduced size and function. With regard to HIV infection, the finding that all affected individuals have circulating anti-CD43 antibodies has led to the suggestion that these auto-antibodies contribute to severe immunodeficiency (Giordanengo *et al.*, (1995) *Blood*, 86:2302-2311).

CD43 is a large, abundant and highly charged trans-membrane molecule that is
30 normally expressed on the surface of white blood cells. In this normal context of

- 2 -

expression, CD43 acts as a barrier molecule preventing intercellular interactions and allowing blood cells to remain in the circulation.

CD43 is composed of 381 amino acids divided between a 235 residue extracellular region, a 23 residue transmembrane region and a 123 amino acid C-terminal intracellular region (Pallant et al., (1989) *Proceedings of the National Academy of Sciences USA*, 86:1328-1332; Shelley et al., (1989) *Proceedings of the National Academy of Sciences USA*, 86:2819-2823). The extracellular region contains approximately 84 sialylated O-linked carbohydrate units and appears by electron microscopy to be a rod-like structure extending 45 nm from the cell surface (Cyster et al., (1991) *EMBO Journal*, 10:893-902). Comparison of the rat, mouse and human sequences indicates that the intracellular domain has been highly conserved during evolution suggesting a critical function. The intracellular domain anchors CD43 to the cytoskeleton by binding actin, ezrin and moesin (Serrador et al., (1998) *Blood*, 91:4632-4644; Yonemura et al., (1998) *Journal of Cell Biology*, 140:885-895).

When leukocytes are at rest, CD43 maintains their circulation within the blood stream by preventing intercellular adhesion. This function is achieved by virtue of the size and strong negative charge of the extracellular domain (Brown et al., (1981) *Nature*, 289:456-460; Ardman et al., (1992) *Proceedings of the National Academy of Sciences USA*, 89:5001-5005; Manjunath et al., (1993) *Journal of Immunology*, 151:1528-1534; Dragone et al., (1995) *Proceedings of the National Academy of Sciences USA*, 92:626-630). During leukocyte activation CD43 expression is dramatically down-regulated allowing intercellular interactions mediated by molecules such as the $\beta 2$ integrins. Intercellular interactions are also actively facilitated by CD43 which, due to changes in glycosylation, switches from being an anti-adhesion to a pro-adhesion molecule (Carlsson et al., (1986) *Journal of Biological Chemistry*, 261:12779-12786; Piller et al., (1988) *Journal of Biological Chemistry*, 263:15146-15150; Remold-O'Donnell et al., (1990) *Journal of Immunology*, 145:3372-3378. 22. Campanero et al., (1991) *European Journal of Immunology*, 21:3045-3048; Bazil et al., (1993) *Proceedings of the National Academy of Sciences USA*, 90:3792-3796; Ellies et al., (1994) *Glycobiology*, 4:885-893; Remold-O'Donnell et al. (1994) *Journal of Immunology*, 152:3595-3605; Tomlinson Jones et al., (1994) *Journal of Immunology*, 153:3426-3439;

Ellies et al., (1996) *Blood*, 88:1725-1732; Weber et al., (1997) *Immunobiology*, 197:82-96).

The pro-adhesive function of CD43 is indicated by its identification as a counter receptor for galectin-1, ICAM-1 and the macrophage adhesion molecule sialoadhesin (Rosenstein et al., (1991) *Nature*, 354:233-235; Baum et al., (1995) *Journal of Experimental Medicine*, 181:877-887; Perillo et al., (1995) *Nature*, 378:736-739; Van den Berg et al., (2001) *Journal of Immunology*, 166:3637-3640). In addition, antibodies to CD43 have been shown to activate monocytes, B lymphocytes, dendritic, mast and natural killer cells (Vargas-Cortes et al., (1988) *Scand Journal of Immunology*, 27:661-671; Nong et al., (1989) *Journal of Experimental Medicine*, 170:259-267; Wiken et al., (1989) *Scand Journal of Immunology*, 29:353-361; Wiken et al., (1989) *Scand Journal of Immunology*, 29:363-370; Kuijpers et al., (1992) *Journal of Immunology*, 149:998-1003; Weber et al., (1994) *Immunology*, 82:638-644; Fanales-Belasio et al., (1997) *Ad Exp Med Biology*, 417:207-212). CD43 binds MHC class I molecules and activates T-lymphocytes in a manner independent of both the T-lymphocyte-receptor/CD3 complex and CD28 (Mentzer et al., (1987) *Journal of Experimental Medicine*, 165:1383-1392; Park et al., (1991) *Nature*, 350:706-709; Sperling et al., (1995) *Journal of Experimental Medicine*, 182:139-146; Stöckl et al., (1996) *Journal of Experimental Medicine*, 184:1769-1779). The activation signals of this pathway are mediated through phosphorylation of the intracellular domain of CD43 and its physical interaction with the tyrosine kinases Fyn and Lck and the serine/threonine kinase STANK (Chatila et al., (1988) *Journal of Immunology*, 140:4308-4314; Silverman et al., (1989) *Journal of Immunology*, 142:4194-4200; Pedraza-Alva et al., (1996) *Journal of Biological Chemistry*, 271:27564-27568; Wang et al., (2000) *Cell Immunology*, 205:34-39). Activation signals transduced by CD43 lead to phosphorylation of Shc, induction of the formation of a Shc/Grb2 complex, tyrosine phosphorylation of Vav, mitogen-activated protein kinase activation and nuclear translocation of ERK2 (Pedraza-Alva et al., (1998) *Journal of Biological Chemistry*, 273:14218-14224). Ultimately, these CD43-mediated signals induce the DNA binding activity of the transcription factors AP-1, NF-AT and NF- κ B and activate the genes encoding interleukin-2, CD69 and CD40-L (Santana et al., (2000) *Journal of Biological Chemistry*, 275:31460-31468).

SUMMARY OF THE INVENTION

The invention is based, in part, on the discovery that ovarian tumor cells abnormally express CD43 on their surfaces. The abnormal expression of CD43 on ovarian cells apparently contributes to the development of ovarian tumors. The invention provides methods of diagnosing ovarian tumors and assessing the prognosis and the progression of disease in patients with ovarian tumors. The invention is also based, in part, on the finding that CD43 inhibitors repress the CD43 promoter which mediates progression of tumors and promotes survival or proliferation of white blood cells. The invention provides methods to treat tumors and methods to reduce the number of white blood cells, as well as methods for inhibiting CD promoters.

According to one aspect of the invention, a method for characterizing an ovarian cell is provided. The method comprises determining the presence or absence of a CD43 molecule in an ovarian cell of a subject to characterize the ovarian cell.

A CD43 molecule refers to a CD43 nucleic acid or a CD43 protein. In some embodiments, the CD43 nucleic acid comprises the nucleic acid sequence of SEQ ID NO: 1 (GenBank Accession NO: NM_003123), SEQ ID NO: 3 (GenBank Accession NO: BC012350), SEQ ID NO: 5 (GenBank Accession NO: X60502), SEQ ID NO: 7 (GenBank Accession NO: J04168), SEQ ID NO: 9 (GenBank Accession NO: M61827), SEQ ID NO: 11, (GenBank Accession NO: J04536), or SEQ ID NO: 13 (GenBank Accession NO: X52075), unique fragments and complements of the foregoing. In some embodiments a CD43 protein comprises the protein sequence of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14, unique fragments, and functional variants of the foregoing.

In some embodiments, the method for characterizing an ovarian cell is a diagnostic method wherein the presence of a CD43 molecule in the ovarian cell indicates that the subject has an ovarian tumor.

In some embodiments, the method for characterizing an ovarian cell is a prognostic method wherein the presence of a CD43 molecule in the ovarian cell indicates that the subject has a poor prognosis. A prognosis predicts the probable outcome of a disease or

- 5 -

the prospect of recovery from disease. A poor prognosis, suggests that the outcome of the disease in the subject is poor or the subject has a low chance of recovery from the disease.

In some embodiments, the method for characterizing an ovarian cell is performed *in vivo*. In other embodiments, the method for characterizing an ovarian cell is performed
5 *in vitro*.

Methods of detecting the presence of a CD43 molecule may be done in the presence of a CD43 binding molecule. One example of a CD43 binding molecule is a CD43 antibody.

In some embodiments, the CD43 antibody is bound to a label. In certain
10 embodiments the label is selected from the group consisting of a fluorescent label, an enzyme label, a radioactive label, a nuclear magnetic resonance active label, a luminescent label, and a chromophore label.

A CD43 antibody may be a CD43 polyclonal antibody or a CD43 monoclonal antibody. One example of a CD43 monoclonal antibody is BS1. Other examples of CD43
15 monoclonal antibodies include but are not limited to: MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

According to another aspect of the invention, a method of treating a subject having
20 or at risk of having a tumor is provided. The method comprises administering to a subject in need of such a treatment a CD43 inhibitor in an effective amount to treat the tumor.

In some embodiments, the CD43 inhibitor is a CD43 nucleic acid binding molecule. One example of a CD43 nucleic acid binding molecule is a heterogeneous nuclear protein K (hnRNP-K) molecule. A hnRNP-K molecule refers to a hnRNP-K
25 nucleic acid or to a hnRNP-K protein. In some embodiments the hnRNP-K nucleic acid comprises the nucleic acid sequence of SEQ ID NO: 15, unique fragments and complements of the foregoing. In other embodiments a hnRNP-K comprises the protein sequence of SEQ ID NO: 16, unique fragments, and functional variants of the foregoing.

In some embodiments, the CD43 inhibitor is a transcription factor. In one
30 embodiment, the transcription factor is a Pur α molecule. A Pur α molecule refers to a Pur α nucleic acid or to a Pur α protein. In some embodiments the Pur α nucleic acid

- 6 -

comprises the nucleic acid sequence of SEQ ID NO: 17, unique fragments and complements of the foregoing. In other embodiments a Pur α comprises the protein sequence of SEQ ID NO: 18, unique fragments, and functional variants of the foregoing.

In some embodiments, the CD43 inhibitor is a CD43 antisense molecule.

5 In some embodiments, the CD43 inhibitor is a CD43 antibody. In some embodiments, the CD43 antibody is bound to a radioisotope. Some radioisotopes could emit α radiations. Others could emit β radiations. Still other radioisotopes could emit γ radiations. Examples of radioisotopes that could be used in this invention include: ^{225}Ac , ^{211}At , ^{212}Bi , ^{213}Bi , ^{186}Rh , ^{188}Rh , ^{177}Lu , ^{90}Y , ^{131}I or ^{67}Cu , ^{125}I , ^{123}I or ^{77}Br .

10 In some embodiments, the CD43 antibody is bound to a therapeutic moiety. Examples of therapeutic moieties that may be bound to the CD43 antibody include but are not limited to drugs, toxins or fragments thereof, or enzymes or fragments thereof.

In some embodiments, the drug is a cytotoxic drug. Examples of the cytotoxic drugs include but are not limited to: calicheamicin, esperamicin, methotrexate, doxorubicin, melphalan, chlorambucil, ARA-C, vindesine, mitomycin C, cis-platinum, etoposide, bleomycin, and 5-fluorouracil.

15 CD43 antibodies may be polyclonal CD43 antibodies or monoclonal CD43 antibodies. Examples of monoclonal CD43 antibodies include but are not limited to BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

20 Tumors encompassed by the invention include but are not limited to benign and malignant solid tumors and benign and malignant non-solid tumors. Examples of solid tumors include but are not limited to: biliary tract cancer, brain cancer (including glioblastomas and medulloblastomas), breast cancer, cervical cancer, choriocarcinoma, colon cancer, endometrial cancer, esophageal cancer, gastric cancer, intraepithelial neoplasms, including Bowen's disease and Paget's disease, liver cancer, lung cancer, lymphomas, including Hodgkin's disease and lymphocytic lymphomas, neuroblastomas, oral cancer, including squamous cell carcinoma, ovarian cancer, including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells, pancreatic cancer, prostate cancer, rectal cancer, renal cancer including adenocarcinoma and Wilms tumor,

- 7 -

sarcomas, including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma and osteosarcoma, skin cancer, including melanoma, Kaposi's sarcoma, basocellular cancer and squamous cell cancer, testicular cancer, including germinal tumors (seminomas, and non-seminomas such as teratomas and choriocarcinomas), stromal tumors and germ cell tumors, and thyroid cancer, including thyroid adenocarcinoma and medullary carcinoma.

Examples of non-solid tumors include but are not limited to hematological neoplasms. A hematologic neoplasm includes, for example, lymphoid disorders, myeloid disorders, and AIDS associated leukemias.

Lymphoid disorders include but are not limited to acute lymphocytic leukemia and chronic lymphoproliferative disorders (e.g., lymphomas, myelomas, and chronic lymphoid leukemias). Lymphomas include Hodgkin's disease and non-Hodgkin's lymphoma. Chronic lymphoid leukemias include T cell chronic lymphoid leukemias and B cell chronic lymphoid leukemias.

Myeloid disorders include chronic myeloid disorders such as, for instance, chronic myeloproliferative disorders, myelodysplastic syndrome and acute myeloid leukemia. Chronic myeloproliferative disorders include but are not limited to angiogenic myeloid metaplasia, essential thrombocythemia, chronic myelogenous leukemia, polycythemia vera, and atypical myeloproliferative disorders. Atypical myeloproliferative disorders include, for example, atypical chronic myelogenous leukemia (CML), chronic neutrophilic leukemia, mast cell disease, and chronic eosinophilic leukemia.

A subject includes a mammal, such as a human, non-human primate, cow, horse, pig, sheep, goat, dog, cat, or rodent. In some preferred embodiments, the subject is a human.

In certain embodiments, the method of treatment further comprises administering one or more anti-tumor therapies. Antitumor therapies include surgery, radiation therapy, or chemotherapeutic agents

In one embodiment the tumor is an ovarian cancer. Ovarian tumors include serous cystoma, mucinous cystoma, endometrioid tumor, mesonephric tumor, transitional cell (Brenner) tumor, dermoid tumor, teratoma, fibroma, thecoma, luteoma, granulosa cell tumor, struma ovarii, dysgerminoma, malignant mixed mesodermal tumor, and carcinoma.

- 8 -

Examples of serous cystomas include serous benign cystadenomas, serous cystadenomas, and serous cystadenocarcinomas.

Examples of mucinous cystomas include mucinous benign cystadenomas, mucinous cystadenomas, and mucinous cystadenocarcinomas.

5 Examples of endometrioid tumors include endometrioid benign cysts and endometrioid adenocarcinomas.

Examples of mesonephric tumors include benign mesonephric tumors and mesonephric cystadenocarcinomas.

In some embodiments, the carcinoma is a clear cell carcinoma.

10 In some embodiments, the ovarian tumor has metastasized beyond the ovary.

In some embodiments of this invention treatment of ovarian tumor further comprises an anti-ovarian tumor therapy. Anti-ovarian tumor therapies include anti-ovarian tumor chemotherapeutic agents. Examples of anti-ovarian tumor chemotherapeutic agents include cisplatin (Platinol) or platinum-containing drug combinations. Other anti-ovarian tumor chemotherapeutic agents include one or more
15 agents selected from the group consisting of : vinblastin, bleomycin, etoposide, hexamethylmelanine, gemcitabine, topotecan, isofosfamide, alkylating agents, progestational agents, and antiestrogens.

Examples of the alkylating agents include: Mechlorethamine (nitrogen mustard;
20 HN2; Mustargen), Chlorambucil (Leukeran), Cyclophosphamide (Cytosan), Melphalan (Alkeran), Thiopeta (triethylenethiophosphoramide), Busulfan (Myleran), Carmustine (BCNU), Lomustine (CCNU), Semustine (methyl-CCNU), Altretamine (hexamethylmelanine), Procarbazine (Matulane), Dacarbazine, and Carboplatin (Paraplatin).

Progestational agents include: progesterone, progesterone derivative, 17-ethinyl
25 testosterone derivative, and 19-nortestosterone derivative.

Progesteron derivatives include: hydroxyprogesterone caproate, medroxyprogesterone acetate, and megestrol acetate.

One Example of 17-ethinyl testosterone derivative is Dimethisterone.

Examples of 19-nortestosterone derivatives include: Desogestrel, Norethynodrel,
30 Lynestrenol, Norethindrone, Norethindrone acetate, Ethynodiol diacetate, and L-Norgestrel.

- 9 -

Antiestrogens include: Tamoxifen, anastrozole, and a combination of aminoglutethimide and hydrocortisone.

According to another aspect of the invention, a method for assessing the regression or progression of an ovarian tumor in a subject is provided. The method comprises the steps of measuring in a first ovarian cell obtained from the subject, the presence of a CD43 molecule, measuring in a second ovarian cell obtained from the subject the presence of a CD43 molecule, comparing the presence of the CD43 molecule in the first ovarian cell and the second ovarian cell, wherein a decrease in the presence of the CD43 molecule in the second ovarian cell compared to the first ovarian cell indicates regression of the ovarian tumor, and wherein an increase in the presence of the CD43 molecule in the second ovarian cell compared to the first ovarian cell indicates progression of the ovarian tumor.

In some embodiments, the CD43 molecule is a CD43 nucleic acid. In other embodiments, the CD43 molecule is a CD43 protein.

Measuring the presence of a CD43 molecule may be performed *in vivo* or *in vitro*. In some embodiments, measuring the presence of a CD43 molecule is done in the presence of a CD43 binding molecule. Examples of CD43 binding molecules are listed above.

According to yet another aspect of the invention, a kit for diagnosing an ovarian tumor is provided. The kit comprises one or more CD43 binding molecules, one or more control agents, and instructions for the use of the CD43 binding molecules, and the control agents in the diagnosis of an ovarian tumor.

In some embodiments, the CD43 binding molecule binds a CD43 nucleic acid. In other embodiments, the CD43 binding molecule binds a CD43 protein.

The kit may be used for diagnosing an ovarian tumor *in vivo* or *in vitro*. CD43 binding molecules that may be used in this aspect of the invention include the CD43 binding molecules described above.

In some embodiments, one or more control agents of the kit are bound to a substrate.

In still another aspect of the invention, a method of reducing the number of white blood cells in a subject is provided. The method involves administering to a subject in

- 10 -

need of a reduction of the number of white blood cells, a CD43 inhibitor molecule in an effective amount to reduce the number of white blood cells in the subject.

In some embodiments, the CD43 inhibitor is a hnRNP-K molecule. In some embodiments, the CD43 inhibitor is a transcription factor such as a Pur α molecule. In some embodiments, the CD43 inhibitor is a CD43 antisense molecule. Examples of CD43 antisense molecules are described above. In some embodiments, the CD43 inhibitor is a CD43 antibody. CD43 antibodies may be polyclonal CD43 antibodies or monoclonal CD43 antibodies. Examples of monoclonal CD43 antibodies are listed above.

Examples of subjects in need of a reduction in the number of white blood cells include subjects with white blood cell disorders such as acute leukemias, chronic leukemias, and lymphomas, and subjects who are about to undergo, are undergoing, or have undergone a bone marrow, organ, cellular, or material transplant.

In some embodiments, the method involves administering a CD43 inhibitor with a second therapy to reduce the number of white blood cells.

In yet another aspect of the invention, a method of inhibiting a CD promoter is provided. The method involves contacting the CD promoter with a CD43 inhibitor in an amount effective to inhibit the CD promoter.

CD43 inhibitors are described above.

In some embodiments of the invention the CD promoter is inhibited *in vivo*. In other embodiments the CD promoter is inhibited *in vitro*.

Examples of CD promoters inhibited by a CD43 inhibitor include but are not limited to: CD43, CD11a, CD11b, CD11c, and CD11d. In some embodiments CD11c is specifically excluded.

These and other aspects of the invention, as well as various advantages and utilities, will be more apparent in reference to the following detailed description of the invention. Each of the limitations of the invention can encompass various embodiments of the invention. It is therefore, anticipated that each of the limitations involving any one element or combination of elements can be included in each aspect of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows down-regulation of CD43 mRNA during K562 activation.

- 11 -

Figure 2 depicts repression of the CD43 promoter during K562 activation.

Figure 3 demonstrates *in vitro* binding of recombinant hnRNP-K binds to the +18/+39 region of the CD43 promoter.

Figure 4 depicts determination of the molecular mass of the Jurkat nuclear factor
5 which binds the +18/+39 region of the CD43 promoter.

Figure 5 shows repression by hnRNP-K of the CD43 promoter present within an extrachromosomal plasmid.

Figure 6 shows repression by hnRNP-K of the CD43 promoter present within chromosomal DNA.

10 Figure 7 is a Southern blot analysis of the transcript produced from the CD43 promoter linked to luciferase and integrated within the K562 genome.

Figure 8 shows that hnRNP-K and Pura α act together to repress the CD43 promoter.

Figure 9 demonstrates a putative secondary structure of the CD43 promoter.

Figure 10 shows the detection of CD43 expression in ovarian cancer specimens by
15 immunocytochemistry.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based, in part, on the discovery that CD43 plays a role in the diagnosis and treatment of tumors. One aspect of the present invention relates to a
20 method of characterizing an ovarian cell based on determining the presence or absence of a CD43 molecule in the ovarian cell obtained from a subject wherein the presence of the CD43 molecule indicates that the subject has an ovarian tumor.

Methods for determining the presence or absence of a CD43 molecule in an ovarian cell are provided. The invention is directed to determining the presence or absence of a
25 CD43 molecule in a ovarian cell and not on a non-ovarian cell that has metastasized to the ovary. In some embodiments, the method of determining the presence or absence of a CD43 molecule is performed in the presence of a CD43 binding molecule. This method involves providing a CD43 binding molecule which binds to a CD43 molecule of an ovarian cell. The cell is contacted with the CD43 binding molecule under conditions
30 effective to permit binding of the CD43 binding molecule to the CD43 molecule of the cell or portions thereof. The CD43 binding molecule may be bound to a label that permits the

- 12 -

detection of the ovarian cell or portions thereof upon binding of CD43 binding molecule to the ovarian cell or portions thereof. The presence of a CD43 molecule is detected by the detection of the label.

In some embodiments, the contact between the CD43 binding molecule and the ovarian cell is carried out *in vivo* in a living mammal and involves administering the CD43 binding molecule to the mammal under conditions that permit binding of the CD43 binding molecule to a CD43 molecule. Such administration can be carried out by any suitable method known to one of ordinary skill in the art.

In other embodiments, the contact between the CD43 binding molecule and the ovarian cell is carried out *in vitro* and involves contacting the CD43 binding molecule with an ovarian cell obtained from a mammal under conditions that permit binding of the CD43 binding molecule to a CD43 molecule. Such contacting can be carried out by any suitable method known to one of ordinary skill in the art. Other methods involve isolation of a cellular fraction(s), i.e., proteins or nucleic acids from the cell and contacting the fraction with a CD43 binding molecule.

As used herein, a "CD43 binding molecule" is a molecule that selectively binds to a CD43 molecule. The CD43 binding molecule binds to a CD43 nucleic acid, and /or unique fragments thereof, or to a CD43 protein and /or unique fragments thereof. Such CD43 binding molecules may be used to selectively bind to a CD43 molecule in a cell (*in vivo* or *ex vivo*) for imaging and therapeutic applications in which, for example, the binding molecule is tagged with a detectable label and/or a toxin for targeted delivery to the CD43 molecule.

In some embodiments, the CD43 binding molecule is a CD43 binding peptide or a CD43 antibody, e.g., an Fab or F(ab)₂ fragment of a CD43 antibody, or an antigen binding fragment of a CD43 antibody. Typically, the antigen binding fragment includes a CDR3 region that is selective for the CD43 molecule. Any of the various types of antibodies can be used for this purpose, including, for example, monoclonal antibodies, polyclonal antibodies, humanized antibodies and chimeric antibodies.

Thus, the invention provides agents which bind to CD43 molecules, and in certain embodiments to unique fragments of the CD43 molecule. Such binding partners can be used in screening assays to determine the presence or absence of a CD43 molecule and in

- 13 -

purification protocols to isolate such CD43 molecules. Likewise, such binding partners can be used to selectively target drugs, toxins or other molecules to cells which express CD43 molecules. In this manner, cells present in solid or non-solid tumors which express CD43 molecules can be treated with cytotoxic compounds. Such agents also can be used
5 to inhibit the native activity of the CD43 molecules, for example, by binding to such polypeptides, to further characterize the functions of these molecules

Thus, a CD43 antibody is an antibody or fragment thereof having the ability to selectively bind to CD43 molecules, and to unique fragments thereof. Antibodies include polyclonal, monoclonal, and chimeric antibodies, prepared, for example, according to
10 conventional methodology. Examples of CD43 antibodies include but are not limited to BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

Other CD43 antibodies may be prepared by any of a variety of methods, including
15 administering protein, fragments of protein, cells expressing the protein or fragments thereof and the like to an animal to induce polyclonal antibodies. The production of monoclonal antibodies is according to techniques well known in the art. Such antibodies may be used, for example, to identify tissues expressing protein or to purify protein. Antibodies also may be coupled to specific labeling agents for imaging or to antitumor
20 agents, including, for example, methotrexate, radioiodinated compounds, toxins such as ricin, other cytostatic or cytolytic drugs, and so forth.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) *The Experimental Foundations of Modern Immunology* Wiley
25 & Sons, Inc., New York; Roitt, I. (1991) *Essential Immunology*, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')₂ fragment, retains both of the antigen binding sites of an intact
30 antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment,

- 14 -

retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains
5 without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

The term "antigen-binding fragment" of an antibody as used herein, refers to one or more portions of an antibody that retain the ability to specifically bind to an antigen (e.g., CD43 molecule). It has been shown that the antigen-binding function of an antibody can
10 be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding fragment" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the V_L , V_H , C_L and C_{H1} domains; (ii) a $F(ab')_2$ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the V_H and C_{H1} domains; (iv)
15 a Fv fragment consisting of the V_L and V_H domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546) which consists of a V_H domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, V_L and V_H , are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made
20 as a single protein chain in which the V_L and V_H regions pair to form monovalent molecules (known as single chain Fv (scFv); see e.g., Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. These antibody fragments are obtained using conventional
25 procedures, such as proteolytic fragmentation procedures, as described in J. Goding, *Monoclonal Antibodies: Principles and Practice*, pp 98-118 (N.Y. Academic Press 1983), which is hereby incorporated by reference as well as by other techniques known to those with skill in the art. The fragments are screened for utility in the same manner as are intact antibodies.

30 Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the

- 15 -

epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, W.R. (1986) *The Experimental Foundations of Modern Immunology* Wiley & Sons, Inc., New York; Roitt, I. (1991) *Essential Immunology*, 7th Ed., Blackwell Scientific Publications, Oxford). In both the heavy chain
5 Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

The non-CDR regions of a mammalian antibody may be replaced with similar
10 regions of nonspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of
15 humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the methods of the
20 invention involve the use of F(ab')₂, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')₂ fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab
25 fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies. Thus, the invention involves polypeptides of
30 numerous size and type that bind specifically to a CD43 molecule. These polypeptides may be derived also from sources other than antibody technology. For example, such

- 16 -

polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries.

Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptides and non-peptide synthetic moieties.

5 The antibodies useful in the methods of the invention encompass various antibody isotypes, such as IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD, IgE. As used herein, "isotype" refers to the antibody class (e.g. IgM or IgG1) that is encoded by heavy chain constant region genes. The antibodies can be full length or can include only an antigen-binding fragment such as the antibody constant and/or variable domain of IgG1,
10 IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgAsec, IgD or IgE or could consist of a Fab fragment, a F(ab')₂ fragment, and a Fv fragment.

 The antibodies useful according to the present invention can be polyclonal, monoclonal, or a mixture of polyclonal and monoclonal antibodies. The antibodies can be produced by a variety of techniques well known in the art. Procedures for raising
15 polyclonal antibodies are well known. For example, anti-CD43 polyclonal antibodies may be raised by administering a CD43 molecule subcutaneously to New Zealand white rabbits which have first been bled to obtain pre-immune serum. The CD43 molecule can be injected at a total volume of 100 µl per site at six different sites, typically with one or more adjustments. The rabbits are then bled two weeks after the first injection and periodically
20 boosted with the same antigen three times every six weeks. A sample of serum is collected 10 days after each boost. Polyclonal antibodies are recovered from the serum, preferably by affinity chromatography using a CD43 molecule to capture the antibody. This and other procedures for raising polyclonal antibodies are disclosed in E. Harlow, et. al., editors, *Antibodies: A Laboratory Manual* (1988), which is hereby incorporated by
25 reference.

 Monoclonal antibody production may be effected by techniques which are also well known in the art. The term "monoclonal antibody," as used herein, refers to a preparation of antibody molecules of single molecular composition. A monoclonal antibody displays a single binding specificity and affinity for a particular epitope. The
30 process of monoclonal antibody production involves obtaining immune somatic cells with the potential for producing antibody, in particular B lymphocytes, which have been

- 17 -

previously immunized with the antigen of interest either *in vivo* or *in vitro* and that are suitable for fusion with a B-cell myeloma line.

An antibody can be linked to a detectable marker, an antitumor agent or an immunomodulator. Antitumor agents can include cytotoxic agents and agents that act on tumor neovasculature. Detectable markers include, for example, radioactive or fluorescent markers. Cytotoxic agents include cytotoxic radionuclides, chemical toxins or chemotherapeutic agents and protein toxins.

The cytotoxic radionuclide or radiotherapeutic isotope may be an alpha-emitting isotope such as ^{225}Ac , ^{211}At , ^{212}Bi , or ^{213}Bi . Alternatively, the cytotoxic radionuclide may a beta-emitting isotope such as ^{186}Rh , ^{188}Rh , ^{177}Lu , ^{90}Y , ^{131}I or ^{67}Cu . Further, the cytotoxic radionuclide may emit low energy electrons and include the isotopes ^{125}I , ^{123}I or ^{77}Br .

Suitable chemical toxins or chemotherapeutic agents include, for example, members of the enediyne family of molecules, such as calicheamicin and esperamicin. Chemical toxins can also be taken from the group consisting of methotrexate, doxorubicin, melphalan, chlorambucil, ARA-C, vindesine, mitomycin C, cis-platinum, etoposide, bleomycin and 5-fluorouracil. Of course, combinations of the various toxins could also be coupled to one antibody molecule thereby accommodating variable cytotoxicity. Other chemotherapeutic agents are known to those skilled in the art.

The coupling of one or more toxin molecules to the CD43 antibody is envisioned to include many chemical mechanisms, for instance covalent binding, affinity binding, intercalation, coordinate binding, and complexation. The toxic compounds used to prepare the anti-CD43 immunotoxins are attached to the CD43 antibodies thereof by standard protocols known in the art.

The covalent binding can be achieved either by direct condensation of existing side chains or by the incorporation of external bridging molecules. Many bivalent or polyvalent agents are useful in coupling protein molecules to other proteins, peptides or amine functions, etc. For example, the literature is replete with coupling agents such as carbodiimides, diisocyanates, glutaraldehyde, diazobenzenes, and hexamethylene diamines. This list is not intended to be exhaustive of the various coupling agents known in the art but, rather, is exemplary of the more common coupling agents.

- 18 -

In some embodiments, it is contemplated that one may wish to first derivatize the antibody, and then attach the toxin component to the derivatized product. Suitable cross-linking agents for use in this manner include, for example, SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), and SMPT, 4-succinimidyl-oxycarbonyl-methyl-(2-pyridyldithio)toluene.

In addition, protein toxins can be fused to the CD43 antibody by genetic methods to form a hybrid immunotoxin fusion protein. To make a fusion immunotoxin protein in accordance with the invention, a nucleic acid molecule is generated that encodes a CD43 antibody, a fragment of CD43 antibody, a single chain CD43 antibody, or a subunit of a CD43 antibody linked to a protein toxin. Such fusion proteins contain at least a targeting agent (e.g., CD43 antibody subunit) and a toxin, operatively attached. The fusion proteins may also include additional peptide sequences, such as peptide spacers which operatively attach the targeting agent and toxin compound, as long as such additional sequences do not appreciably affect the targeting or toxin activities of the fusion protein. The two proteins can be attached by a peptide linker or spacer, such as a glycine-serine spacer peptide, or a peptide hinge, as is well known in the art. Thus, for example, the C-terminus of a CD43 antibody can be fused to the N-terminus of the protein toxin molecule to form an immunotoxin that retains the binding properties of the CD43 antibody. Other fusion arrangements will be known to one of ordinary skill in the art.

To express the fusion immunotoxin, the nucleic acid encoding the fusion protein is inserted into an expression vector in accordance with standard methods, for stable expression of the fusion protein, preferably in mammalian cells, such as CHO cells. The fusion protein can be isolated and purified from the cells or culture supernatant using standard methodology, such as a CD43 affinity column.

Radionuclides typically are coupled to an antibody by chelation. For example, in the case of metallic radionuclides, a bifunctional chelator is commonly used to link the isotope to the antibody or other protein of interest. Typically, the chelator is first attached to the antibody, and the chelator-antibody conjugate is contacted with the metallic radioisotope. A number of bifunctional chelators have been developed for this purpose, including the diethylenetriamine pentaacetic acid (DTPA) series of amino acids described in U.S. patents 5,124,471, 5,286,850 and 5,434,287, which are incorporated herein by

- 19 -

reference. As another example, hydroxamic acid-based bifunctional chelating agents are described in U.S. patent 5,756,825, the contents of which are incorporated herein. Another example is the chelating agent termed *p*-SCN-Bz-HEHA (1,4,7,10,13,16-hexaazacyclo-octadecane- N,N',N'',N''',N'''',N'''''-hexaacetic acid) (Deal et al., *J. Med. Chem.* 42:2988, 5 1999), which is an effective chelator of radiometals such as ²²⁵Ac.

In addition, the CD43 antibodies can be used in immunofluorescence techniques to examine human tissue, cell and bodily fluid specimens. In one protocol, slides containing cryostat sections of frozen, unfixed tissue biopsy samples or cytological smears are air dried, formalin or acetone fixed, and incubated with the monoclonal antibody preparation 10 in a humidified chamber at room temperature. The slides are then washed and further incubated with a preparation of a secondary antibody directed against the monoclonal antibody, usually some type of anti-mouse immunoglobulin if the monoclonal antibodies used are derived from the fusion of a mouse spleen lymphocyte and a mouse myeloma cell line. This secondary antibody is tagged with a compound, for instance rhodamine or 15 fluorescein isothiocyanate, that fluoresces at a particular wavelength. The staining pattern and intensities within the sample are then determined by fluorescent light microscopy and optionally photographically recorded.

As another alternative, computer enhanced fluorescence image analysis or flow cytometry can be used to examine tissue specimens or cells, i.e., single cell preparations 20 from aspiration biopsies of tumors using CD43 antibodies. The CD43 antibodies are particularly useful in quantitation of live tumor cells, i.e., single cell preparations from biopsies of tumors by computer enhanced fluorescence image analyzer or with a flow cytometer. The antibodies are particularly useful, for example, in such assays to differentiate benign from malignant ovarian tumors since the CD43 molecule to which the 25 CD43 antibody binds is expressed in increased amounts by malignant tumors as compared to benign ovarian tumors. The percent CD43 positive cell population, alone or in conjunction with determination of other attributes of the cells (e.g., DNA ploidy of these cells), may, additionally, provide very useful prognostic information by providing an early indicator of disease progression.

30 The antibodies may also be used in combination with other known antibodies to provide additional information regarding the malignant phenotype of a tumor.

- 20 -

The method of the present invention can be used to screen patients for diseases associated with the presence of tumors. Alternatively, it can be used to identify the recurrence of such diseases, particularly when the disease is localized in a particular biological material of the patient. Using the method of the present invention, this
5 recurrence can be detected by administering a short range radiolabeled antibody to the mammal and then detecting the label.

Alternatively, the contacting step can be carried out in a biological sample. When the contacting is carried out in a biological sample, it is preferred that the biological sample recognize substantially no antigens other than CD43. Thus, the biological samples
10 and methods of the present invention can be used to determine the effectiveness of a tumor treatment protocol by monitoring the level of CD43 in a biological sample.

The CD43 antibodies or antigen-binding fragments thereof can also be utilized in *in vivo* therapy of tumors. As used herein, therapy includes prophylaxis, which encompasses preventing or delaying the development or the progression of a tumor.

15 The antibodies can be used alone or covalently attached, either directly or via linker, to a compound which kills and/or inhibits proliferation of the tumor following administration and localization of the conjugates. When the antibody is used by itself, it may mediate tumor destruction by complement fixation or antibody-dependent cellular cytotoxicity. Alternatively, the antibody may be administered in combination with a
20 chemotherapeutic drug to cause synergistic therapeutic effects (Baslya and Mendelsohn, 1994 *Breast Cancer Res. and Treatment* 29:127-138). A variety of different types of substances can be directly conjugated to the antibody for therapeutic uses, including radioactive metal and non-metal isotopes, chemotherapeutic drugs, toxins, etc. as described above and known in the art (see, e.g., Vitetta and Uhr, 1985, *Annu. Rev.*
25 *Immunol.* 3:197).

The CD43 antibodies thereof can also be administered together with complement. Accordingly, within the scope of the invention are compositions comprising antibodies or antigen-binding fragments thereof and serum or complement. These compositions are advantageous in that the complement is located in close proximity to the human antibodies
30 or antigen-binding fragments thereof. Alternatively, the antibodies or antigen-binding

- 21 -

fragments thereof of the invention and the complement or serum can be administered separately.

The CD43 antibodies can be administered with one or more immunostimulatory agents to induce or enhance an immune response, such as IL-2 and immunostimulatory oligonucleotides (e.g., those containing CpG motifs). Preferred immunostimulatory agents stimulate specific arms of the immune system, such as natural killer (NK) cells that mediate antibody-dependent cell cytotoxicity (ADCC).

Antigens, such as the CD43 molecule, can be administered with one or more adjuvants to induce or enhance an immune response. An adjuvant is a substance which potentiates the immune response. Adjuvants of many kinds are well known in the art. Specific examples of adjuvants include monophosphoryl lipid A (MPL, SmithKline Beecham); saponins including QS21 (SmithKline Beecham); immunostimulatory oligonucleotides (e.g., CpG oligonucleotides described by Kreig et al., *Nature* 374:546-9, 1995); incomplete Freund's adjuvant; complete Freund's adjuvant; montanide; vitamin E and various water-in-oil emulsions prepared from biodegradable oils such as squalene and/or tocopherol.

Other agents which stimulate the immune response of the subject to a CD43 molecule can also be administered to the subject. For example, cytokines are also useful in vaccination protocols as a result of their lymphocyte regulatory properties. Many cytokines useful for such purposes will be known to one of ordinary skill in the art, including interleukin-2 (IL-2); IL-12, which has been shown to enhance the protective effects of vaccines (*see, e.g., Science* 268: 1432-1434, 1995); GM-CSF; IL-18 and the like. Thus, cytokines can be administered in conjunction with antibodies, antigens, and/or adjuvants to increase an immune response.

The antibodies or antigen-binding fragments thereof can be used in conjunction with other therapeutic treatment modalities. Such other treatments include surgery, radiation, cryosurgery, thermotherapy, hormone treatment, chemotherapy, vaccines, and other immunotherapies.

Also within the scope of the invention are kits to measure the levels of a CD43 molecule and instructions for use. The kits could be used for detecting (diagnosing, prognosis, or monitoring) a tumor. The kits can further contain a least one additional

- 22 -

reagent, such as complement, or one or more additional antibodies of the invention (e.g., an antibody having a complementary activity which binds to an epitope in the CD43 molecule distinct from the first antibody).

Kits containing the CD43 antibodies can be prepared for *in vitro* diagnosis, prognosis and/or monitoring a tumor by the immunohistological, immunocytological and immunoserological methods described above. The components of the kits can be packaged either in aqueous medium or in lyophilized form. When the antibodies are used in the kits in the form of conjugates in which a label moiety is attached, such as an enzyme or a radioactive metal ion, the components of such conjugates can be supplied either in fully conjugated form, in the form of intermediates or as separate moieties to be conjugated by the user of the kit.

A kit may comprise a carrier being compartmentalized to receive in close confinement therein one or more container means or series of container means such as test tubes, vials, flasks, bottles, syringes, or the like. A first of said container means or series of container means may contain one or more CD43 antibodies or a CD43 molecule. A second container means or series of container means may contain a label or linker-label intermediate capable of binding to the primary CD43 antibody (or fragment thereof).

Kits for use in *in vivo* tumor localization and therapy method containing the CD43 antibodies thereof conjugated to other compounds or substances can be prepared. The components of the kits can be packaged either in aqueous medium or in lyophilized form. When the antibodies or antigen-binding fragments thereof are used in the kits in the form of conjugates in which a label or a therapeutic moiety is attached, such as a radioactive metal ion or a therapeutic drug moiety, the components of such conjugates can be supplied either in fully conjugated form, in the form of intermediates or as separate moieties to be conjugated by the user of the kit.

It should be understood that kits could be assembled as to provide convenient access and use in clinical settings. Individuals skilled in the art can readily modify packaging to suit individual needs.

The invention in some aspects involves a method of treating a subject having or at risk of having a tumor. The method involves administering to a subject in need of such treatment a CD43 inhibitor in an effective amount to treat the tumor.

- 23 -

As used herein, a "CD43 inhibitor" is a molecule that inhibits (reduces or eliminates) the expression of a CD43 molecule or inhibits the function of a CD43 molecule.

As used herein, a "CD43 molecule" refers to a CD43 nucleic acid or to a CD43 protein. A CD43 nucleic acid refers to a nucleic acid which encodes a "CD43 protein." In some embodiments the CD43 nucleic acid is selected from the group consisting of:

(a) nucleic acids which hybridize under stringent conditions to a nucleic acid having a nucleotide sequence set forth as SEQ ID NO: 1 (GenBank Accession NO: NM_003123), SEQ ID NO: 3 (GenBank Accession NO: BC012350), SEQ ID NO: 5 (GenBank Accession NO: X60502), SEQ ID NO: 7 (GenBank Accession NO: J04168), SEQ ID NO: 9 (GenBank Accession NO: M61827), SEQ ID NO: 11, (GenBank Accession NO: J04536), or SEQ ID NO: 13 (GenBank Accession NO: X52075) and which code for a CD43 protein,

(b) deletions, additions and substitutions of the nucleic acids of (a), and

(c) nucleic acids that differ from the nucleic acids of (a) or (b) in codon sequence due to the degeneracy of the genetic code.

As used herein a "CD43 protein" refers to a protein having the activity of a native CD43 protein such as that described by SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14, as well as unique fragments, functionally active fragments or variants thereof, provided that the functional fragments or variants exhibit a CD43 functional activity. The term protein as used herein encompasses the terms polypeptide and peptide. As used herein, a CD43 functional activity refers, for example, to the ability of a CD43 protein to prevent intercellular adhesions.

The invention embraces variants of the CD43 proteins described herein. As used herein, a "variant" of a CD43 protein is a protein which contains one or more modifications to the primary amino acid sequence of a CD43 protein. Modifications which create a CD43 protein variant can be made to a CD43 protein 1) to produce, increase, reduce, or eliminate activity of the CD43 protein; 2) to enhance a property of the CD43 protein, such as protein stability in an expression system or the stability of protein-protein binding; or 3) to provide a novel activity or property to a CD43 protein.

- 24 -

Modifications to a CD43 protein are typically made to the CD43 nucleic acid which encodes the CD43 protein, and can include deletions, point mutations, truncations, amino acid substitutions and additions of amino acids or non-amino acid moieties. Alternatively, modifications can be made directly to the protein, such as by cleavage, addition of a linker molecule, addition of a detectable moiety, such as biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the CD43 amino acid sequences.

One of skill in the art will be familiar with methods for predicting the effect on protein conformation of a change in protein sequence, and can thus "design" a variant protein (e.g. CD43) according to known methods. One example of such a method is described by Dahiyat and Mayo in *Science* 278:82-87, 1997, whereby proteins can be designed *de novo*. The method can be applied to a known protein to vary only a portion of the protein sequence. By applying the computational methods of Dahiyat and Mayo, specific variants of a protein can be proposed and tested to determine whether the variant retains a desired conformation.

In general, functional variants of a protein include proteins which are modified specifically to alter a feature of the protein unrelated to its desired physiological activity. The skilled artisan will also realize that conservative amino acid substitutions may be made in the protein to provide functional variants of the foregoing proteins, i.e., the variants which have the functional capabilities of the protein. As used herein, a "conservative amino acid substitution" refers to an amino acid substitution which does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

Such substitutions can be made by a variety of methods known to one of ordinary skill in the art. For example, amino acid substitutions may be made by PCR-directed mutation, site-directed mutagenesis according to the method of Kunkel (*Proc. Nat. Acad. Sci. U.S.A.* 82: 488-492, 1985), or by chemical synthesis of a gene encoding a protein. Where amino acid substitutions are made to a small unique fragment of a protein the substitutions can be made by directly synthesizing the protein. The activity of functional

- 25 -

variants or fragments of protein can be tested by cloning the gene encoding the altered protein into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the altered protein, and testing for a functional capability of the protein as disclosed herein.

5 Variants of a protein can be prepared according to methods for altering protein sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John
10 Wiley & Sons, Inc., New York.

In general, the treatment methods involve administering a CD43 inhibitor to treat a tumor. In some embodiments, the CD43 inhibitor is a CD43 binding molecule. One example of a CD43 binding molecule is a hnRNP-K molecule.

As used herein, a "hnRNP-K molecule" refers to a hnRNP-K nucleic acid or to a
15 hnRNP-K protein. A hnRNP-K nucleic acid refers to a nucleic acid which encodes a "hnRNP protein" and complements thereof. In some embodiments the hnRNP nucleic acid is selected from the group consisting of:

(a) nucleic acids which hybridize under stringent conditions to a nucleic acid having a nucleotide sequence set forth as SEQ ID NO:15 (GenBank Accession NO:
20 NM_031263.1) and which code for a hnRNP-K protein,

(b) deletions, additions and substitutions of the nucleic acids of (a),

(c) nucleic acids that differ from the nucleic acids of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and

(d) complements of (a), (b), or (c).

25 One preferred hnRNP nucleic acid is an hnRNP-K nucleic acid which encodes a hnRNP-K protein having the amino acid sequence of SEQ ID NO: 16 (GenBank Accession NO: AAB20770.1).

Another example of a CD43 binding molecule is a Pur α molecule. It has been discovered that the combination of an hnRNP molecule and a Pur α molecule produce an
30 even greater physiological response than either molecule alone. As used herein, a "Pur α molecule" refers to a Pur α nucleic acid or to a Pur α protein. A Pur α nucleic acid refers to

- 26 -

a nucleic acid which encodes a Pura protein. This includes, for instance, nucleic acids selected from the group consisting of:

- (a) nucleic acids which hybridize under stringent conditions to a nucleic acid having a nucleotide sequence set forth as SEQ ID NO: 17 (GenBank Accession NO: M96684.1) and which code for a Pura protein,
- (b) deletions, additions and substitutions of the nucleic acids of (a),
- (c) nucleic acids that differ from the nucleic acids of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and
- (d) complements of (a), (b), or (c).

One preferred Pura nucleic acid is one which encodes a Pura protein having an amino acid sequence of SEQ ID NO: 18 (GenBank Accession NO: NM_005859.2).

The nucleic acids of the invention may be DNA, RNA, mixed deoxyribonucleotides and ribonucleotides, or may also incorporate synthetic non-natural nucleotides.

The term "stringent conditions" as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. More specifically, stringent conditions, as used herein, refers, for example, to hybridization at 65°C in hybridization buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% Bovine Serum Albumin, 2.5mM NaH₂PO₄ (pH 7), 0.5% SDS, 2mM EDTA). SSC is 0.15M sodium chloride/0.15M sodium citrate, pH 7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid.

After hybridization, the membrane upon which the DNA is transferred is washed at 2 x SSC at room temperature and then at 0.1 x SSC/0.1 x SDS at temperatures up to 68°C.

The foregoing set of hybridization conditions is but one example of stringent hybridization conditions known to one of ordinary skill in the art. There are other conditions, reagents, and so forth which can be used, which result in a stringent hybridization. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to

- 27 -

manipulate the conditions in a manner to permit the clear identification of homologs and alleles of the nucleic acids of the invention. The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules which then are routinely isolated, followed by isolation of the pertinent nucleic acid and sequencing.

5 In general CD43 homologs and alleles typically will share at least 40% nucleotide identity to SEQ ID NO: 1 (GenBank Accession NO: NM_003123), SEQ ID NO: 3 (GenBank Accession NO: BC012350), SEQ ID NO: 5 (GenBank Accession NO: X60502), SEQ ID NO: 7 (GenBank Accession NO: J04168), SEQ ID NO: 9 (GenBank Accession NO: M61827), SEQ ID NO: 11 (GenBank Accession NO: J04536), or SEQ ID
10 NO: 13 (GenBank Accession NO: X52075) and/or at least 50% amino acid identity to NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14, in some instances will share at least 50% nucleotide identity and/or at least 65% amino acid identity and in still other instances will share at least 60% nucleotide identity and/or at least 75% amino acid identity. Preferred homologs and alleles share
15 nucleotide and amino acid identities with SEQ ID NO: 1 (GenBank Accession NO: NM_003123), SEQ ID NO: 3 (GenBank Accession NO: BC012350), SEQ ID NO: 5 (GenBank Accession NO: X60502), SEQ ID NO: 7 (GenBank Accession NO: J04168), SEQ ID NO: 9 (GenBank Accession NO: M61827), SEQ ID NO: 11, (GenBank Accession NO: J04536), or SEQ ID NO: 13 (GenBank Accession NO: X52075) and SEQ ID NO: 2,
20 SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, or SEQ ID NO: 14, respectively, and encode polypeptides of greater than 80%, more preferably greater than 90%, still more preferably greater than 95% and most preferably greater than 99% identity.

In general, hnRNP-K homologs and alleles typically will share at least 40%
25 nucleotide identity and/or at least 50% amino acid identity to SEQ ID NO: 15 and SEQ ID NO: 16, respectively, in some instances will share at least 50% nucleotide identity and/or at least 65% amino acid identity and in still other instances will share at least 60% nucleotide identity and/or at least 75% amino acid identity. Preferred homologs and alleles share nucleotide and amino acid identities with SEQ ID NO: 15 and SEQ ID NO: 16 ,
30 respectively, and encode polypeptides of greater than 80%, more preferably greater than 90%, still more preferably greater than 95% and most preferably greater than 99% identity.

- 28 -

Pur α homologs and alleles typically will share at least 40% nucleotide identity and/or at least 50% amino acid identity to SEQ ID NO: 17 and SEQ ID NO: 18, respectively, in some instances will share at least 50% nucleotide identity and/or at least 65% amino acid identity and in still other instances will share at least 60% nucleotide
5 identity and/or at least 75% amino acid identity. Preferred homologs and alleles share nucleotide and amino acid identities with SEQ ID NO: 17 and SEQ ID NO: 18, respectively, and encode polypeptides of greater than 80%, more preferably greater than 90%, still more preferably greater than 95% and most preferably greater than 99% identity.

The percent identity can be calculated using various, publicly available software
10 tools developed by NCBI (Bethesda, Maryland) that can be obtained through the internet. Exemplary tools include the BLAST system which uses algorithms developed by Altschul et al. (*Nucleic Acids Res.* 25:3389-3402, 1997). Pairwise and ClustalW alignments (BLOSUM30 matrix setting) as well as Kyte-Doolittle hydropathic analysis can be obtained using the MacVector sequence analysis software (Oxford Molecular Group).
15 Watson-Crick complements of the foregoing nucleic acid molecules also are embraced by the invention.

The invention also includes degenerate nucleic acids which include alternative codons to those present in the native materials. For example, serine residues are encoded by the codons TCA, AGT, TCC, TCG, TCT and AGC. Each of the six codons is
20 equivalent for the purposes of encoding a serine residue. Thus, it will be apparent to one of ordinary skill in the art that any of the serine-encoding nucleotide triplets may be employed to direct the protein synthesis apparatus, *in vitro* or *in vivo*, to incorporate a serine residue into an elongating protein such as a CD43 protein, a hnRNP-K protein, or a Pur α protein. Similarly, nucleotide sequence triplets which encode other amino acid
25 residues include, but are not limited to: CCA, CCC, CCG and CCT (proline codons); CGA, CGC, CGG, CGT, AGA and AGG (arginine codons); ACA, ACC, ACG and ACT (threonine codons); AAC and AAT (asparagine codons); and ATA, ATC and ATT (isoleucine codons). Other amino acid residues may be encoded similarly by multiple nucleotide sequences. Thus, the invention embraces degenerate nucleic acids that differ
30 from the biologically isolated nucleic acids in codon sequence due to the degeneracy of the genetic code.

- 29 -

The nucleic acids useful herein may be isolated nucleic acids. As used herein, the term "isolated nucleic" acid means: (i) amplified *in vitro* by, for example, polymerase chain reaction (PCR); (ii) recombinantly produced by cloning; (iii) purified, as by cleavage and gel separation; or (iv) synthesized by, for example, chemical synthesis. An isolated nucleic acid is one which is readily manipulable by recombinant DNA techniques well known in the art. Thus, a nucleotide sequence contained in a vector in which 5' and 3' restriction sites are known or for which polymerase chain reaction (PCR) primer sequences have been disclosed is considered isolated but a nucleic acid sequence existing in its native state in its natural host is not. An isolated nucleic acid may be substantially purified, but need not be. For example, a nucleic acid that is isolated within a cloning or expression vector is not pure in that it may comprise only a tiny percentage of the material in the cell in which it resides. Such a nucleic acid is isolated, however, as the term is used herein because it is readily manipulable by standard techniques known to those of ordinary skill in the art. An isolated nucleic acid as used herein encompasses a non-naturally occurring chromosome as well as total genomic DNA isolated from cells and tissues.

In some aspects of the invention, the treatment methods involve administering to a subject a nucleic acid (e.g. hnRNP-K nucleic acid and/or a Pur α nucleic acid) to treat a tumor. In some embodiments, the nucleic acid is introduced *ex vivo* into the cell(s) of a subject and the cell(s) are returned to the subject. The nucleic acid is under operable control of regulatory elements which permit expression of the nucleic acid in the cell(s). Numerous transfection and transduction techniques as well as appropriate expression vectors are well known to those of ordinary skill in the art, some of which are described in PCT application WO95/00654. *In vivo* introduction of the hnRNP-K nucleic acid or Pur α nucleic acid using vectors such as adenovirus, retroviruses, herpes virus, and targeted liposomes is also contemplated according to the invention.

In preferred embodiments, a virus vector for delivering a nucleic acid encoding a protein of the invention (e.g. hnRNP-protein or Pur α protein) is selected from the group consisting of adenoviruses, adeno-associated viruses, poxviruses including vaccinia viruses and attenuated poxviruses, Semliki Forest virus, Venezuelan equine encephalitis virus, retroviruses, Sindbis virus, and Ty virus-like particle. Examples of viruses and virus-like particles which have been used to deliver exogenous nucleic acids include:

- 30 -

replication-defective adenoviruses (e.g., Xiang et al., *Virology* 219:220-227, 1996; Elo et al., *J. Virol.* 71:5375-5381, 1997; Chengalvala et al., *Vaccine* 15:335-339, 1997), a modified retrovirus (Townsend et al., *J. Virol.* 71:3365-3374, 1997), a nonreplicating retrovirus (Irwin et al., *J. Virol.* 68:5036-5044, 1994), a replication defective Semliki Forest virus (Zhao et al., *Proc. Natl. Acad. Sci. USA* 92:3009-3013, 1995), canarypox virus and highly attenuated vaccinia virus derivative (Paoletti, *Proc. Natl. Acad. Sci. USA* 93:11349-11353, 1996), non-replicative vaccinia virus (Moss, *Proc. Natl. Acad. Sci. USA* 93:11341-11348, 1996), replicative vaccinia virus (Moss, *Dev. Biol. Stand.* 82:55-63, 1994), Venezuelan equine encephalitis virus (Davis et al., *J. Virol.* 70:3781-3787, 1996), Sindbis virus (Pugachev et al., *Virology* 212:587-594, 1995), and Ty virus-like particle (Allsopp et al., *Eur. J. Immunol* 26:1951-1959, 1996). In one of the preferred embodiments, the virus vector is an adenovirus.

Another one of the preferred viruses for certain applications is the adeno-associated virus, a double-stranded DNA virus. The adeno-associated virus is capable of infecting a wide range of cell types and species and can be engineered to be replication-deficient. It further has advantages, such as heat and lipid solvent stability, high transduction frequencies in cells of diverse lineages, including hematopoietic cells, and lack of superinfection inhibition thus allowing multiple series of transductions. In addition, wild-type adeno-associated virus infections have been followed in tissue culture for greater than 100 passages in the absence of selective pressure, implying that the adeno-associated virus genomic integration is a relatively stable event. The adeno-associated virus can also function in an extrachromosomal fashion.

In general, other preferred viral vectors are based on non-cytopathic eukaryotic viruses in which non-essential nucleic acids have been replaced with the nucleic acids of interest. Non-cytopathic viruses include retroviruses, the life cycle of which involves reverse transcription of genomic viral RNA into DNA with subsequent proviral integration into host cellular DNA. In general, the retroviruses are replication-deficient (i.e., capable of directing synthesis of the desired proteins, but incapable of manufacturing an infectious particle). Such genetically altered retroviral expression vectors have general utility for the high-efficiency transduction of nucleic acids *in vivo*. Standard protocols for producing replication-deficient retroviruses (including the steps of incorporation of exogenous

- 31 -

genetic material into a plasmid, transfection of a packaging cell lined with plasmid, production of recombinant retroviruses by the packaging cell line, collection of viral particles from tissue culture media, and infection of the target cells with viral particles) are provided in Kriegler, M., *Gene Transfer and Expression, A Laboratory Manual*, W.H.

5 Freeman Co., New York (1990) and Murry, E.J. Ed. *Methods in Molecular Biology*, vol. 7, Humana Press, Inc., Clifton, New Jersey (1991).

In addition to the viral vectors described above, various techniques may be employed for introducing nucleic acid molecules of the invention into cells, depending on whether the nucleic acid molecules are introduced *in vitro* or *in vivo* in a host. Such techniques include transfection of nucleic acid molecule- CaPO_4 precipitates, transfection of nucleic acid molecules associated with DEAE, transfection or infection with the foregoing viruses including the nucleic acid molecule of interest, liposome mediated transfection, and the like. For certain uses, it is preferred to target the nucleic acid molecule to particular cells. In such instances, a vehicle used for delivering a nucleic acid molecule of the invention into a cell (e.g., a retrovirus, or other virus; a liposome) can have a targeting molecule attached thereto. For example, a molecule such as an antibody specific for a surface membrane protein on the target cell or a ligand for a receptor on the target cell can be bound to or incorporated within the nucleic acid molecule delivery vehicle. In some embodiments, monoclonal antibodies are preferred. Where liposomes are employed to deliver the nucleic acid molecules of the invention, proteins which bind to a surface membrane protein associated with endocytosis may be incorporated into the liposome formulation for targeting and/or to facilitate uptake. Such proteins include capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half life, and the like. Polymeric delivery systems also have been used successfully to deliver nucleic acid molecules into cells, as is known by those skilled in the art. Such systems even permit oral delivery of nucleic acid molecules.

In some other aspects of the invention the treatment methods involve administering to a subject a hnRNP-K protein. The term protein, as used herein encompasses the terms polypeptide and peptide. As used herein a "hnRNP-K protein" refers to a protein having the activity of a native hnRNP-K protein such as that described by SEQ ID NO: 16 as well

- 32 -

as functionally active fragments or variants thereof. Thus, for example a hnRNP-K protein includes proteins which include the proteins of SEQ ID NO: 16, unique fragments, and functional variants thereof, as well as proteins which are encoded by nucleic acids such as those having SEQ ID NO: 15, provided that the functional fragment or variant exhibits a
5 hnRNP-K functional activity. As used herein, a hnRNP-K functional activity refers to the ability of a hnRNP-K protein to modulate one or more of the following parameters such as: suppressing tumor growth and/or cell proliferation, reducing the number of white blood cells, and repressing CD molecule expression. An exemplary hnRNP-K functional activity is a tumor suppressor activity such as suppressing and/or reducing tumor cell growth,
10 proliferation, and/or metastasis.

A hnRNP-K functional activity can be determined, for example, by assaying tumor size in a biological sample. An exemplary assay involves: measuring the tumor size in a biological sample before treatment with a putative hnRNP-K molecule, measuring the tumor size in the biological sample after treatment with the putative hnRNP-K molecule,
15 and comparing the tumor size in the biological sample before and after treatment with the putative hnRNP-K molecule. In this example, a decrease or even less of an increase than would be observed in the absence of such treatment, in the tumor size in the biological sample after treatment with the putative hnRNP-K molecule indicates that the putative hnRNP-K molecule is an hnRNP-K having a hnRNP-K functional activity. An increase in
20 the tumor size in the biological sample after treatment with the putative hnRNP-K molecule over that which would occur in the absence of a hnRNP-K molecule indicates that the putative hnRNP-K molecule is not a hnRNP molecule of the invention and does not have a hnRNP-K functional activity.

The hnRNP-K functional activity can be determined *ex vivo* or *in vitro*. An *ex vivo*
25 or *in vitro* hnRNP-K functional activity assay involves, for example, assaying the number of tumor cells in an *ex vivo* medium or in an *in vitro* sample. The assay involves: measuring the number of tumor cells in an *ex vivo* medium or *in vitro* sample before treatment with a putative hnRNP-K molecule, measuring the number of tumor cells in the *ex vivo* medium or *in vitro* sample after treatment with the putative hnRNP-K molecule,
30 and comparing the number of tumor cells in the *ex vivo* medium or *in vitro* sample before and after treatment with the putative hnRNP-K molecule. A decrease or less of an increase

- 33 -

(compared to a control) in the number of tumor cells in the *ex vivo* medium or *in vitro* sample after treatment with the putative hnRNP-K molecule indicates that the putative hnRNP-K molecule has a hnRNP-K functional activity. A suitable control is a known hnRNP-K protein such as that identified by SEQ ID NO: 16. An increase in the number of tumor cells in comparison with a control in the *ex vivo* medium or *in vitro* sample after treatment with the putative hnRNP-K molecule indicates that the putative hnRNP-K molecule is not a hnRNP-K molecule of the invention and does not have a hnRNP-K functional activity.

Although not wishing to be bound to any particular theory or mechanism, it is believed that the hnRNP-K protein may affect at least some of the above-noted cell functions by inhibiting a CD promoter. Thus *in vitro* assays analyzing CD promoter activation with putative hnRNP-K molecules may also be useful for identifying hnRNP-K molecules having hnRNP-K functional activity.

The treatment methods may optionally involve administering to a subject a Pur α protein. As used herein a Pur α protein refers to a protein having the activity of a native Pur α protein such as that described by SEQ ID NO: 18 as well as functionally active fragments or variants thereof. Thus, for example a Pur α protein includes proteins which include the proteins of SEQ ID NO: 18, unique fragments, and functional variants thereof, as well as proteins which are encoded by nucleic acids such as those having SEQ ID NO: 17, provided that the functional fragment or variant exhibits a Pur α functional activity. As used herein, a Pur α functional activity refers to the ability of a Pur α protein to modulate one or more of the following parameters such as: suppressing tumor growth and/or cell proliferation, reducing the number of white blood cells, and repressing CD molecule expression. An exemplary Pur α functional activity is a tumor suppressor activity such as suppressing and/or reducing tumor cell growth, proliferation, and/or metastasis. A Pur α functional activity can be determined using any of the exemplary assays set forth above for hnRNP-K molecules.

The hnRNP-K proteins are isolated proteins. Proteins can be isolated from biological samples including tissue or cell homogenates, and can also be expressed recombinantly in a variety of prokaryotic and eukaryotic expression systems by constructing an expression vector appropriate to the expression system, introducing the

- 34 -

expression vector into the expression system, and isolating the recombinantly expressed protein. Short proteins, also can be synthesized chemically using well-established methods of peptide synthesis.

Thus, as used herein with respect to proteins, "isolated" means separated from its native environment and present in sufficient quantity to permit its identification or use. Isolated, when referring to a protein means, for example: (i) selectively produced by expression of a recombinant nucleic acid or (ii) purified as by chromatography or electrophoresis. Isolated proteins may, but need not be, substantially pure. The term "substantially pure" means that the proteins are essentially free of other substances with which they may be found in nature or *in vivo* systems to an extent practical and appropriate for their intended use. Substantially pure proteins may be produced by techniques well known in the art. Because an isolated protein may be admixed with a pharmaceutically acceptable carrier in a pharmaceutical preparation, the protein may comprise only a small percentage by weight of the preparation. The protein is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, e.g. isolated from other proteins.

A fragment of a protein (CD43, hnRNP-K, or Pur α for example) generally has the features and characteristics of fragments including unique fragments. As will be recognized by those skilled in the art, the size of a fragment which is unique will depend upon factors such as whether the fragment constitutes a portion of a conserved protein domain. Thus, some regions of proteins will require longer segments to be unique while others will require only short segments, typically between 5 and 12 amino acids (e.g. 5, 6, 7, 8, 9, 10, 11, and 12 amino acids long).

Unique fragments of a protein preferably are those fragments which retain a distinct functional capability of the protein. Functional capabilities which can be retained in a fragment of a protein include interaction with antibodies, interaction with other proteins or fragments thereof, selective binding of nucleic acid molecules, and enzymatic activity. One important activity is the ability to act as a signature for identifying the protein.

Those skilled in the art are well versed in methods for selecting unique amino acid sequences, typically on the basis of the ability of the fragment to selectively distinguish the

- 35 -

sequence of interest from non-family members. A comparison of the sequence of the fragment to those on known data bases typically is all that is necessary.

The invention embraces variants of the hnRNP-K proteins described herein. As used herein, a "variant" of a hnRNP-K protein is a protein which contains one or more
5 modifications to the primary amino acid sequence of a hnRNP-K protein. Modifications which create a hnRNP-K protein variant can be made to a hnRNP-K protein 1) to produce, increase, reduce, or eliminate activity of the hnRNP-K protein; 2) to enhance a property of the hnRNP-K protein, such as protein stability in an expression system or the stability of protein-protein binding; 3) to provide a novel activity or property to a hnRNP-K protein;
10 or 4) to provide equivalent or better binding to an CD promoter. Modifications to a hnRNP-K protein are typically made to the hnRNP-K nucleic acid which encodes the hnRNP-K protein, and can include deletions, point mutations, truncations, amino acid substitutions and additions of amino acids or non-amino acid moieties. Alternatively, modifications can be made directly to the protein, such as by cleavage, addition of a linker
15 molecule, addition of a detectable moiety, such as biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the hnRNP-K amino acid sequences.

The invention also embraces variants of the Pura proteins described herein. As used herein, a "variant" of a Pura protein is a protein which contains one or more
20 modifications to the primary amino acid sequence of a Pura protein. Modifications which create a Pura protein variant can be made to a Pura protein 1) to produce, increase, reduce, or eliminate activity of the Pura protein; 2) to enhance a property of the Pura protein, such as protein stability in an expression system or the stability of protein-protein binding; 3) to provide a novel activity or property to a Pura protein; or 4) to provide
25 equivalent or better binding to an CD promoter. Modifications to a Pura protein are typically made to the Pura nucleic acid which encodes the Pura protein, and can include deletions, point mutations, truncations, amino acid substitutions and additions of amino acids or non-amino acid moieties. Alternatively, modifications can be made directly to the protein, such as by cleavage, addition of a linker molecule, addition of a detectable moiety,
30 such as biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the Pura amino acid sequences.

- 36 -

Upon determining that a peptide derived from a hnRNP-K protein plays a role in, for example, suppression of tumor growth and/or cell proliferation, reduction in the number of white blood cells, or repression of CD molecule expression, one can make conservative amino acid substitutions to the amino acid sequence of the peptide to make a
5 variant of a hnRNP-K protein. The substituted peptides can then be tested for one or more of the above-noted functions, *in vivo* or *in vitro*. These variants can be tested for improved stability and are useful, *inter alia*, in pharmaceutical compositions.

Exemplary functional variants of the hnRNP-K protein include conservative amino acid substitutions of proteins encoded by SEQ ID NO: 16. Conservative amino-acid
10 substitutions in the amino acid sequence of the hnRNP-K protein to produce functional variants of hnRNP-K protein typically are made by alteration of the nucleic acid molecule encoding a hnRNP-K protein. Exemplary functional variants of the Pur α protein include conservative amino acid substitutions of proteins encoded by SEQ ID NO: 18.

Conservative amino-acid substitutions in the amino acid sequence of the Pur α protein to
15 produce functional variants of a Pur α protein typically are made by alteration of the nucleic acid molecule encoding a Pur α protein.

The invention also embraces CD43 antisense molecules that selectively bind to a nucleic acid molecule encoding a CD43 nucleic acid, unique fragments, or complements of these nucleic acid molecules. A unique fragment is one that is a 'signature' for the larger
20 nucleic acid. It, for example, is long enough to assure that its precise sequence is not found in molecules outside of the defined nucleic acid molecules. Those of ordinary skill in the art may apply no more than routine procedures to determine if a fragment is unique within the genome.

As used herein, the term "antisense oligonucleotide" or "antisense" describes an
25 oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that mRNA. The antisense molecules are designed so as to interfere with transcription or
30 translation of a target gene upon hybridization with the target gene or transcript. Those skilled in the art will recognize that the exact length of the antisense oligonucleotide and

- 37 -

its degree of complementarity with its target will depend upon the specific target selected, including the sequence of the target and the particular bases which comprise that sequence.

It is preferred that the antisense oligonucleotide be constructed and arranged so as to bind selectively with the target under physiological conditions, i.e., to hybridize

5 substantially more to the target sequence than to any other sequence in the target cell under physiological conditions. Based upon CD43 antisense molecules, the known sequences for CD43 or upon allelic or homologous genomic and/or cDNA sequences, one of skill in the art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. In order to be sufficiently selective and
10 potent for inhibition, such antisense oligonucleotides should comprise at least 10 and, more preferably, at least 15 consecutive bases which are -complementary to the target, although in certain cases modified oligonucleotides as short as 7 bases in length have been used successfully as antisense oligonucleotides (Wagner et al., Nat. Med.

1(11):1116-1118, 1995). Even more preferably, the antisense oligonucleotides comprise a
15 complementary sequence of 20-35 bases. Although oligonucleotides may be chosen which are antisense to any region of the gene (such as to the CD43) or mRNA transcripts, the antisense oligonucleotides preferably correspond to N-terminal or 5' upstream sites such as translation initiation, transcription initiation or promoter sites. In addition, 3'-untranslated regions may be targeted by antisense oligonucleotides. Targeting to mRNA splicing sites
20 has also been used in the art but may be less preferred if alternative mRNA splicing occurs. In addition, the antisense is targeted, preferably, to sites in which mRNA secondary structure is not expected (see, e.g., Sainio et al., *Cell Mol. Neurobiol.* 14(5):439-457, 1994) and at which proteins are not expected to bind.

In some embodiments, the antisense oligonucleotides of the invention may be
25 composed of "natural" deoxyribonucleotides, ribonucleotides, or any combination thereof. That is, the 5' end of one native nucleotide and the 3' end of another native nucleotide may be covalently linked, as in natural systems, via a phosphodiester internucleoside linkage. These oligonucleotides may be prepared by art recognized methods which may be carried out manually or by an automated synthesizer. They also may be produced recombinantly
30 by vectors.

- 38 -

In some embodiments, however, the antisense oligonucleotides of the invention also may include "modified" oligonucleotides. That is, the oligonucleotides may be modified in a number of ways which do not prevent them from hybridizing to their target but which enhance their stability or targeting or which otherwise enhance their therapeutic effectiveness.

The term "modified oligonucleotide" as used herein describes an oligonucleotide in which (1) at least two of its nucleotides are covalently linked via a synthetic internucleoside linkage (i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide) and/or (2) a chemical group not normally associated with nucleic acids has been covalently attached to the oligonucleotide. Preferred synthetic internucleoside linkages are phosphorothioates, alkylphosphonates, phosphorodithioates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters and peptides.

The term "modified oligonucleotide" also encompasses oligonucleotides with a covalently modified base and/or sugar. For example, modified oligonucleotides include oligonucleotides having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified oligonucleotides may include a 2'-O-alkylated ribose group. In addition, modified oligonucleotides may include sugars such as arabinose instead of ribose.

Delivery systems for molecules of the invention can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the CD43 inhibitor increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di- and tri-glycerides;

- 39 -

hydrogel release systems; sylastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which an agent of the invention is contained in a form within a matrix such as those
5 described in U.S. Patent Nos. 4,452,775, 4,675,189, and 5,736,152, and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation.

Use of a long-term sustained release implant may be particularly suitable for
10 treatment of chronic conditions. Long-term release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term sustained release implants are well-known to those of ordinary skill in the art and include some of the release systems described above.

As used herein, a subject is a mammal such as a human, non-human primate, cow,
15 horse, pig, sheep, goat, dog, cat, or rodent. In some preferred embodiments, the subject is a human. Exemplary subjects calling for treatment with a CD43 inhibitor include subjects having or at risk of having a tumor (solid and non-solid). In some embodiments a subject is free of disorders previously known to be treated using a CD43 inhibitor.

A subject having a tumor is a subject with at least one identifiable sign, symptom,
20 or laboratory finding sufficient to make a diagnosis of a tumor in accordance with clinical standards known in the art for identifying such tumors. Examples of such clinical standards can be found in Harrison's Principles of Internal Medicine, 15th Ed., Fauci AS et al., eds., McGraw-Hill, New York, 2001. In some instances, a diagnosis of a tumor will include identification of a particular cell type present in a sample of a body fluid or tissue
25 obtained from the subject.

A subject at risk of having a tumor is a subject with an identifiable risk factor for having a tumor. For example, a subject at risk of having a tumor can include an individual with a known or suspected exposure to environmental agents (e.g., carcinogens) associated with an increased risk of having a tumor. Additionally or alternatively, a subject at risk of
30 having a tumor can include an individual with a genetic predisposition to developing a tumor. Yet other examples of a subject at risk of having a tumor include a subject that

- 40 -

previously has been diagnosed with a cancer associated with a tumor and who is at risk of metastasis of the primary tumor.

As used herein, "treating" includes preventing, delaying, abating or arresting the clinical symptoms and or signs of a tumor. Treatment also includes reducing or preventing tumor cell growth, proliferation, and/or metastasis as well as increasing the resistance of a subject to develop a disease.

Tumors encompassed by the invention include, for example, benign and malignant solid tumors and benign and malignant non-solid tumors. Examples of solid tumors include but are not limited to: biliary tract cancer, brain cancer (including glioblastomas and medulloblastomas), breast cancer, cervical cancer, choriocarcinoma, colon cancer, endometrial cancer, esophageal cancer, gastric cancer, intraepithelial neoplasms, including Bowen's disease and Paget's disease, liver cancer, lung cancer, lymphomas, including Hodgkin's disease and lymphocytic lymphomas, neuroblastomas, oral cancer, including squamous cell carcinoma, ovarian cancer, including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells, pancreatic cancer, prostate cancer, rectal cancer, renal cancer including adenocarcinoma and Wilms tumor, sarcomas, including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma and osteosarcoma, skin cancer, including melanoma, Kaposi's sarcoma, basocellular cancer and squamous cell cancer, testicular cancer, including germinal tumors (seminomas, and non-seminomas such as teratomas and choriocarcinomas), stromal tumors and germ cell tumors, and thyroid cancer, including thyroid adenocarcinoma and medullary carcinoma.

Examples of non-solid tumors include but are not limited to hematological neoplasms. As used herein, a hematologic neoplasm is a term of art which includes lymphoid disorders, myeloid disorders, and AIDS associated leukemias.

Lymphoid disorders include but are not limited to acute lymphocytic leukemia and chronic lymphoproliferative disorders (e.g., lymphomas, myelomas, and chronic lymphoid leukemias). Lymphomas include, for example, Hodgkin's disease and non-Hodgkin's lymphoma. Chronic lymphoid leukemias include, for example, T cell chronic lymphoid leukemias and B cell chronic lymphoid leukemias.

Myeloid disorders include chronic myeloid disorders such as for instance chronic myeloproliferative disorders and myelodysplastic syndrome and acute myeloid leukemia.

- 41 -

Chronic myeloproliferative disorders include but are not limited to angiogenic myeloid metaplasia, essential thrombocythemia, chronic myelogenous leukemia, polycythemia vera, and atypical myeloproliferative disorders. Atypical myeloproliferative disorders include atypical chronic myelogenous leukemia, chronic neutrophilic leukemia, mast cell
5 disease, and chronic eosinophilic leukemia.

The CD43 inhibitor may be administered alone or in conjunction with one or more therapies known or believed to be useful for treating a tumor. Such therapies include, for example, anti-tumor therapy such as radiation therapy, surgery, or a chemotherapeutic agent or a combination thereof. As used herein, an anti-tumor therapy refers to a therapy
10 that is administered to a subject for the purpose of treating a tumor. Various types of therapies for the treatment of tumors are described herein. Anti-tumor therapies include immunotherapeutic agents, tumor vaccines, hormone therapy, and biological response modifiers.

Anti-tumor therapies function in a variety of ways. For example, some anti-tumor
15 therapies work by targeting physiological mechanisms that are specific to tumor cells. Examples include the targeting of specific genes and their gene products (i.e., proteins primarily) which are mutated in tumors. Such genes include but are not limited to oncogenes (e.g., Ras, Her2, bcl-2), tumor suppressor genes (e.g., EGF, p53, Rb), and cell cycle targets (e.g., CDK4, p21, telomerase). Some anti-tumor therapies can alternately
20 target signal transduction pathways and molecular mechanisms which are altered in tumor cells. Targeting of tumor cells via the epitopes expressed on their cell surface is accomplished through the use of monoclonal antibodies. This latter type of anti-tumor therapy is generally referred to herein as immunotherapy.

Other anti-tumor therapies target cells other than tumor cells. For example, some
25 anti-tumor therapies prime the immune system to attack tumor cells (i.e., tumor vaccines). Still other anti-tumor therapies, called angiogenesis inhibitors, function by attacking the blood supply of solid tumors. Since the most malignant tumors are able to metastasize (i.e., exist the primary tumor site and seed a distal tissue, thereby forming a secondary tumor), therapies that impede this metastasis are also useful in the treatment of tumor.
30 Angiogenic mediators include basic FGF, VEGF, angiopoietins, angiostatin, endostatin, TNF- α , TNP-470, thrombospondin-1, platelet factor 4, CAI, and certain members of the

- 42 -

integrin family of proteins. One category of this type of anti-tumor therapy is a metalloproteinase inhibitor, which inhibits the enzymes used by the tumor cells to exit the primary tumor site and extravasate into another tissue.

Immunotherapeutic agents are medicaments which derive from antibodies or
5 antibody fragments which specifically bind or recognize a tumor antigen. As used herein a tumor antigen is broadly defined as an antigen expressed by a tumor cell. In some embodiments, the antigen is expressed at the cell surface of the tumor cell. In some preferred embodiments, the antigen is one which is not expressed by normal cells, or at least not expressed to the same level as in tumor cells. Antibody-based immunotherapies
10 may function by binding to the cell surface of a tumor cell and thereby stimulate the endogenous immune system to attack the tumor cell. Another way in which antibody-based therapy functions is as a delivery system for the specific targeting of toxic substances to tumor cells. Antibodies are usually conjugated to toxins such as ricin (e.g., from castor beans), calicheamicin and maytansinoids, to radioactive isotopes such as
15 Iodine-131 and Yttrium-90, to chemotherapeutic agents (as described herein), or to biological response modifiers. In this way, the toxic substances can be concentrated in the region of the tumor and non-specific toxicity to normal cells can be minimized. In addition to the use of antibodies which are specific for tumor antigens, antibodies which bind to vasculature, such as those which bind to endothelial cells, are also useful in the
20 invention. This is because generally solid tumors are dependent upon newly formed blood vessels to survive, and thus most tumors are capable of recruiting and stimulating the growth of new blood vessels. As a result, one strategy of many tumor medicaments is to attack the blood vessels feeding a tumor and/or the connective tissues (or stroma) supporting such blood vessels.

25 Chemotherapeutic agents which can be used according to the invention include but are not limited to Aminoglutethimide, Asparaginase, Busulfan, Carboplatin, Chlorombucil, Cytarabine HCl, Dactinomycin, Daunorubicin HCl, Estramustine phosphate sodium, Etoposide (VP16-213), Floxuridine, Fluorouracil (5-FU), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alfa-2a, Alfa-2b, Leuprolide acetate (LHRH-releasing factor analogue), Lomustine (CCNU), Mechlorethamine HCl (nitrogen mustard),
30 Mercaptopurine, Mesna, Mitotane (o.p'-DDD), Mitoxantrone HCl, Octreotide, Plicamycin,

Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Amsacrine (m-AMSA), Azacitidine, Erythropoietin, Hexamethylmelamine (HMM), Interleukin 2, Mitoguazone (methyl-GAG; methyl glyoxal bis-guanyldrazone; MGBG), Pentostatin (2'deoxycoformycin), Semustine (methyl-CCNU), Teniposide (VM-26) and Vindesine sulfate.

Tumor vaccines are medicaments which are intended to stimulate an endogenous immune response against tumor cells. Currently produced vaccines predominantly activate the humoral immune system (i.e., the antibody dependent immune response). Other vaccines currently in development are focused on activating the cell-mediated immune system including cytotoxic T lymphocytes which are capable of killing tumor cells. Tumor vaccines generally enhance the presentation of tumor antigens to both antigen presenting cells (e.g., macrophages and dendritic cells) and/or to other immune cells such as T cells, B cells, and NK cells. In some instances, tumor vaccines may be used along with adjuvants, such as those described above.

Tumor antigens, such as those present in tumor vaccines or those used to prepare tumor immunotherapies, can be prepared from crude tumor cell extracts, as described in Cohen PA et al. (1994) *Cancer Res* 54:1055-8, or by partially purifying the antigens, using recombinant technology, or de novo synthesis of known antigens. Tumor antigens can be used in the form of immunogenic portions of a particular antigen or in some instances a whole cell or a tumor mass can be used as the antigen. Such antigens can be isolated or prepared recombinantly or by any other means known in the art.

Other vaccines take the form of dendritic cells which have been exposed to tumor antigens *in vitro*, have processed the antigens and are able to express the tumor antigens at their cell surface in the context of MHC molecules for effective antigen presentation to other immune system cells. Dendritic cells form the link between the innate and the acquired immune system by presenting antigens and through their expression of pattern recognition receptors which detect microbial molecules like LPS in their local environment.

The invention also provides a method of reducing the number of white blood cells in a subject. The method involves administering to a subject in need of such reduction in the number of white blood cells, a CD43 inhibitor in an effective amount to reduce the

- 44 -

number of white blood cells in said subject. Exemplary subjects in need of a reduction of the number of white blood cells include subjects with white blood cell disorders such as acute leukemias, chronic leukemias, and lymphomas. Other examples of subjects requiring reduction in the number of white blood cells include subjects who are about to
5 undergo, are undergoing, or have undergone a bone marrow, organ, cellular, or material transplant. A subject about to undergo a bone marrow, organ, cellular, or material transplant is a subject having a disorder requiring a bone marrow, organ, cellular or material transplant.

As used herein reduction of the number of white bloods refers to decreasing the
10 number of existing white blood cells or decreasing the production of new white blood cells. Decreasing, as used herein, includes total elimination or eradication, as well as other decreases which do not result in total eradication.

A transplant is an organ, tissue, a cell(s), or a material for grafting into an area of the body or into another individual: A transplant encompasses an organ, tissue, or a cell(s)
15 taken from the body for grafting into another area of the same body or into another individual. A transplant also encompasses *ex-vivo* material (organ, tissue, cell(s) or material inert or non-inert, artificial or non-artificial) for grafting into an area of the body. As used herein, *ex-vivo* material refers to material prepared outside the body. Examples of transplants include: allogeneic transplants (transplants of tissue between genetically
20 dissimilar animals of the same species, heterotopic transplants (transplants of tissue typical of one area to a different recipient site), orthotopic or homotypic transplants (transplants of tissue typical of one area to an identical recipient site), syngeneic transplants (transplants of tissues between animals in the same pure line, e.g., within an inbred strain), syngenesioplasmic transplants (transplants of tissue from one individual to a related
25 individual of the same species).

The CD43 inhibitor may be administered alone or in conjunction with one or more therapies known or believed to be useful to reduce the number of white blood cells. Examples of medicaments that reduce the number of white blood cells include for example, cytarabine (cytosine arabinoside, anthracyclines such as daunorubicin, Atr, arsenic trioxide, hydroxyurea, busulphan, homoharringtonine, and interferons (IFN) such
30 as IFN- α , IFN- β , and IFN- γ .

- 45 -

Those skilled in the art will recognize which of the other therapies to be administered in conjunction with the CD43 inhibitor are appropriate for treating a given suspected or identified tumor or reducing the number of white blood cells.

The invention provides pharmaceutical compositions of CD43 inhibitor. The pharmaceutical compositions contains a CD43 inhibitor in a pharmaceutically acceptable carrier. Thus, in a related aspect, the invention provides a method for forming a medicament that involves placing a therapeutically effective amount of the CD43 inhibitor in the pharmaceutically acceptable carrier to form one or more doses.

When administered, the therapeutic compositions of the present invention are administered in pharmaceutically acceptable preparations. Such preparations may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives, compatible carriers, supplementary immune potentiating agents such as adjuvants and cytokines and optionally other therapeutic agents.

The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue, or organism. The characteristics of the carrier will depend on the route of administration. Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art.

An effective amount means, with respect to a tumor for example, that amount of a CD43 inhibitor which prevents the onset of, alleviates the symptoms of, or stops or slows the progression of a tumor or reduces the number of white blood cells. In general such symptoms are, at least in part, the result of unwanted cell proliferation cells *in vivo*. Thus, a tumor is a condition that is characterized by certain clinical features and which, it is generally believed, is associated with unwanted cell proliferation cells *in vivo*.

"Unwanted," with respect to cell proliferation cells *in vivo*, refers to cell proliferation which results in an adverse medical condition. The phrase "therapeutically effective amount" means, with respect to treatment of a tumor or reduction in the number of white blood cells which prevents the onset of, alleviates the symptoms of, or stops the signs of a tumor or reduces the number of white blood cells.

- 46 -

The pharmaceutical preparations disclosed herein are prepared in accordance with standard procedures and are administered at dosages that are selected to reduce, prevent or eliminate the condition (See, e.g., Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA, and Goodman and Gilman's The Pharmaceutical Basis of Therapeutics, Pergamon Press, New York, N.Y., the contents of which are incorporated herein by reference, for a general description of the methods for administering various agents for human therapy).

The pharmaceutically acceptable compositions of the present invention comprise a CD43 inhibitor in association with one or more nontoxic, pharmaceutically acceptable carriers and/or diluents and/or adjuvants and/or excipients, collectively referred to herein as "carrier" materials, and if desired other active ingredients.

The CD43 inhibitor of the present invention may be administered by any route, preferably in the form of a pharmaceutical composition adapted to such a route, and would be dependent on the condition being treated. The compounds and compositions may, for example, be administered orally, intravascularly, intramuscularly, subcutaneously, intraperitoneally, or topically. Preferred routes of administration include oral and intravenous administration.

For oral administration, the CD43 inhibitor may be in the form of, for example, a tablet, capsule, suspension or liquid. The pharmaceutical composition is preferably made in the form of a dosage unit containing a therapeutically effective amount of the active ingredient. Examples of such dosage units are tablets and capsules. For therapeutic purposes, the tablets and capsules can contain, in addition to the active ingredient, conventional carriers such as binding agents, for example, acacia gum, gelatin, polyvinylpyrrolidone, sorbitol, or tragacanth; fillers, for example, calcium phosphate, cellulose, glycine, lactose, maize-starch, mannitol, sorbitol, or sucrose; lubricants, for example, magnesium stearate, polyethylene glycol, silica, or talc; disintegrants, for example potato starch, flavoring or coloring agents, or acceptable wetting agents. Oral liquid preparations generally in the form of aqueous or oily solutions, suspensions, emulsions, syrups or elixirs may contain conventional additives such as suspending agents, emulsifying agents, non-aqueous agents, preservatives, coloring agents and flavoring agents. Examples of additives for liquid preparations include acacia, almond oil, ethyl

- 47 -

alcohol, fractionated coconut oil, gelatin, glucose syrup, glycerin, hydrogenated edible fats, lecithin, methyl cellulose, methyl or propyl para-hydroxybenzoate, propylene glycol, sorbitol, or sorbic acid.

The pharmaceutical compositions may also be administered via injection.

5 Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions or suspensions may be prepared from sterile powders or granules having one or more of the carriers mentioned for use in the formulations for oral administration. The compounds may be dissolved in polyethylene glycol, propylene glycol, ethanol, corn oil, benzyl alcohol, sodium chloride,
10 sterile water, and/or various buffers.

For topical use the compounds of the present invention may also be prepared in suitable forms to be applied to the skin, or mucus membranes of the nose and throat, and may take the form of creams, ointments, liquid sprays or inhalants, lozenges, or throat
15 paints. Such topical formulations further can include chemical compounds such as dimethylsulfoxide (DMSO) to facilitate surface penetration of the active ingredient. Suitable carriers for topical administration include oil-in-water or water-in-oil emulsions using mineral oils, petrolatum and the like, as well as gels such as hydrogel. Alternative
topical formulations include shampoo preparations, oral pastes and mouthwash.

For rectal administration the compounds of the present invention may be
20 administered in the form of suppositories admixed with conventional carriers such as cocoa butter, wax or other glyceride.

Alternatively, the compounds of the present invention may be in powder form for reconstitution at the time of delivery.

The dosage regimen for treating a tumor or for reducing the number of white blood
25 cells with a CD43 inhibitor is selected in accordance with a variety of factors, including the type, age, weight, sex and medical condition of the subject, the severity of the disease, the route and frequency of administration, the renal and hepatic function of the subject, and the particular compound employed. An ordinarily skilled physician or clinician can readily determine and prescribe the effective amount of the drug required to treat a tumor
30 or to reduce the number of white blood cells. In general, dosages are determined in

- 48 -

accordance with standard practice for optimizing the correct dosage for treating a tumor, reducing the number of white blood cells, or inhibiting a CD promoter.

The dosage regimen can be determined, for example, by following the response to the treatment in terms clinical signs. Examples of such clinical signs are well known in the art, and they include for example the pulse, blood pressure, temperature, and respiratory rate. Harrison's Principles of Internal Medicine, 15th Ed., Fauci AS et al., eds., McGraw-Hill, New York, 2001.

Typically dosages of the CD43 inhibitor will be dependent upon the condition to be treated. In general, the active agent concentration will range from between 0.01 mg per kg of body weight per day (mg/kg/day) to about 10.0 mg/kg/day. Alternatively, the dosages of the CD43 inhibitor will range from between 0.01 micromole per kg of body weight per day (μ mole/kg/day) to about 10 μ mole/kg/day. Preferred oral dosages in humans may range from daily total dosages of about 1-1000 mg/day over the effective treatment period. Preferred intravenous dosages in humans may range from daily total dosages of about 1-100 mg/day over the effective treatment period.

In general, doses of radionuclide delivered by the CD43 antibodies of the invention can range from about 0.01 mCi/Kg to about 10 mCi/kg. Preferably the dose of radionuclide ranges from about 0.1 mCi/Kg to about 1.0 mCi/kg. The optimal dose of a given isotope can be determined empirically by simple routine titration experiments well known to one of ordinary skill in the art.

In yet another aspect, the invention provides a method of inhibiting a CD promoter. The method involves contacting the CD promoter with a CD43 inhibitor in an amount effective to inhibit the CD promoter. CD (cluster of differentiation) refers to a specific antigenic marker found on cells such as lymphocytes. A promoter is a nucleotide sequence of DNA to which RNA polymerase binds and initiates transcription. It usually lies upstream of (5' to) a coding sequence. A promoter sequence aligns the RNA polymerase so that transcription will initiate at a specific site. CD promoters include but are not limited to: CD43, CD11a, CD11b, CD11c, and CD11d.

As used herein, "inhibiting a CD promoter" refers to decreasing or slowing the activity of the CD promoter including halting or eliminating the activity of the CD promoter temporarily or permanently. Inhibition of a CD promoter by a CD43 inhibitor

- 49 -

can be determined, for example, by assaying RNA production or expression of a CD molecule in a sample. Such an assay involves: measuring the expression of the CD molecule in a sample before treatment with a CD43 inhibitor, measuring the expression of the CD molecule in the sample after treatment with a CD43 inhibitor, and comparing the expression of the CD molecule before and after treatment with the CD43 inhibitor. A decrease or no increase in the expression of the CD molecule after treatment with the CD43 inhibitor indicates that CD promoter is inhibited by the CD43 inhibitor. An increase in the expression of the CD molecule after treatment with the CD43 inhibitor indicates the CD promoter is not inhibited by the CD43 inhibitor.

In some embodiments of the invention the CD promoter is inhibited in a biological sample *in vivo*. In other embodiments the CD promoter is inhibited in a biological sample *in vitro*. As used herein, inhibition of the CD promoter refers to decreasing the function of the CD promoter including a total elimination of the CD promoter function.

The biological sample having the CD promoter be located *in vivo* or *in vitro*. For example, the biological sample can be a tissue *in vivo*. Alternatively, the biological sample can be located *in vitro* (e.g., a blood sample, tumor biopsy, tissue extract). In a particularly preferred embodiment, the biological sample can be a cell-containing sample, more preferably a sample containing tumor cells or white blood cells. Samples of tissue and/or cells for use in the various methods described herein can be obtained through standard methods. Samples can be surgical samples of any type of tissue or body fluid. Samples can be used directly or processed to facilitate analysis (e.g., paraffin embedding). Exemplary samples include a cell, a cell scraping, a cell extract, a blood sample, a tissue biopsy, including punch biopsy, a tumor biopsy, a bodily fluid, a tissue, or a tissue extract or other methods.

The invention will be more fully understood by reference to the following examples. These examples, however, are merely intended to illustrate the embodiments of the invention and are not to be construed to limit the scope of the invention.

EXAMPLES

Example 1

Introduction

CD43 is an abundant, heavily glycosylated molecule expressed specifically on the surface of leukocytes and platelets. When leukocytes are at rest CD43 acts to prevent both homotypic and heterotypic interactions. However, during leukocyte activation CD43 expression is repressed facilitating the intercellular contact required for chemotaxis, phagocytosis, aggregation, adhesion to endothelium and transendothelial migration. In this way CD43 repression contributes to both innate and acquired immunity. Here we report that a dramatic down-regulation of CD43 mRNA levels occurs during activation of the leukocytic cell line K562. This repression coincides with repression of the transcriptional activity of the CD43 gene promoter.

We investigated the possibility that CD43 repression is mediated by transcriptional events. We have previously reported that transcriptional repression of the CD43 promoter occurs during activation of the monocytic cell line U937 and that this is mediated by the transcription factor Pur α (Shelley *et al.*, (2001) *Br Journal of Haematology*, 115:159-166). Here we report that transcriptional repression of the CD43 promoter also occurs during activation of the pre-erythroid/pre-megakaryocytic cell line K562. We demonstrated that heterogeneous nuclear ribonucleoprotein K (hnRNP-K) alone and in combination with Pur α mediates repression of the CD43 promoter during K562 activation. HnRNP-K and Pur α bind single-stranded DNA.

Materials and Methods

Cells and Culture Media: Jurkat T-lymphocytic cells and K562 pre-erythroid/pre-megakaryocytic cells were obtained from the American Type Culture Collection (ATCC) and grown in RPMI 1640 medium supplemented with 10% fetal calf serum, 2 mM L-glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin. K562 cells were activated by the addition of phorbol 12-myristate 13-acetate (PMA) (Sigma Chemical Co., St Louis, MO) to a final, concentration of 100 ng/ml. K562 cells stably transfected with the -2/+99 CD43 promoter linked to the luciferase reporter gene were selected using 300 μ g/ml of zeocin (Invitrogen, Carlsbad, CA). The DNA methyltransferase inhibitor 5-azacytidine (Sigma, St. Louis, MO) was used at a concentration of 10 μ M.

RNA Isolation and Northern Blot Analysis: Total RNA was isolated from K562 cells by guanidinium isothiocyanate lysis and centrifugation through a cesium chloride

- 51 -

cushion. 20 µg of total RNA were subjected to electrophoresis through 1% agarose/2.2 M formaldehyde gels and subsequently transferred to Hybond N⁺ nylon membranes (Amersham Pharmacia Biotech Inc., Piscataway, NJ). These membranes were then hybridized with radiolabelled probes which specifically interact with the coding region of CD43 mRNA or glyceraldehyde-3-phosphate dehydrogenase (GAPDH) mRNA. The GAPDH and CD43 coding region probes have been described previously (Shelley et al., (1989) *Proceedings of the National Academy of Sciences USA*, 86:2819-2823; Tso et al., (1985) *Nucleic Acids Research*, 13:2485-2502). Probes were labeled with (α³²P)-dCTP using a NonaPrimer Kit (Appligene, Illkirch, France). Following hybridization, membranes were washed and subjected to autoradiography.

Southern Blot Analysis. DNA was subjected to electrophoresis through a 1% agarose gel and transferred to a Hybond N⁺ nylon membrane (Amersham Pharmacia Biotech Inc., Piscataway, NJ). This membrane was then hybridized with the oligonucleotide *LUC-2* which had been radiolabeled at its 5' end and which specifically interacts with the coding region of the firefly luciferase gene. *LUC-2* was labeled using T4 polynucleotide kinase and [γ³²P]ATP. The nucleotide sequence of *LUC-2* is: 5'-ATAGCCTTATGCAGTTGCTCT-3' (SEQ ID NO: 23). Following hybridization, membranes were washed and subjected to autoradiography.

Plasmid Construction: The activity of the CD43 promoter was assessed using the expression vector pATLuc (Shelley et al., (1993) *Proceedings of the National Academy of Sciences USA*, 90:5364-5368) which contains a promoterless firefly luciferase reporter gene. The polymerase chain reaction (PCR) was used to generate a fragment of the CD43 gene representing nucleotides -2 to +99 relative to the most 5' of the two major transcription initiation sites (Shelley et al., (1990) *Biochemical Journal*, 270:569-576). This fragment was then subcloned into the "filled-in" HindIII site of pATLuc to generate p43Wt (Shelley et al., (2001) *Br Journal of Haematology*, 115:159-166; Farokhzad et al., (2000) *Nucleic Acids Research*, 28:2256-2267). The correct orientation and nucleotide sequence of the CD43 promoter within p43Wt was verified by DNA sequencing (Sanger et al., (1977) *Proceedings of the National Academy of Sciences USA*, 74:5463-5467). K562 cells carrying within their genome the -2/+99 region of the CD43 gene linked to the luciferase reporter were produced using the plasmid p43Wt/Zeo. This plasmid was

- 52 -

generated by inserting between the *SalI* and *PstI* sites of p43Wt the *XhoI/PstI* fragment of pCMV/Zeo (Invitrogen, Carlsbad, CA) containing the zeocin resistance gene. Prior to transfection p43Wt/Zeo was linearized by digestion with *PstI* such that the zeocin gene lay downstream and head-to-tail relative to the luciferase gene. The hnRNP-K expression constructs, Full-Length hnRNP-K, GST-RNP-K and the equivalent vectors empty of hnRNP-K coding sequences were kindly provided by David Levens (National Institutes of Health, Bethesda, MD) (Tomonaga *et al.*, (1995) *Journal of Biological Chemistry*, 270:4875-4881). The Pura expression construct, pHAPur1, was kindly provided by Edward Johnson (Mount Sinai School of Medicine, New York) and the empty vector equivalent, pHA, produced by religation following liberation of the Pura sequence by *RsrII* and *EcoRI* digestion.

Reverse-Transcriptase Polymerase Chain Reaction (RT-PCR). Total RNA was prepared from K562 cells carrying within their genome the CD43 promoter fused to the luciferase reporter. A GeneRacer Kit (Invitrogen Life Technologies, Carlsbad, CA) was used to ligate the RNA oligonucleotide *GeneRacer RNA Oligo* to the 5' end specifically of full-length mRNA within the total RNA mixture. This ligated mRNA was then converted to cDNA using reverse-transcriptase and the *GeneRacer Oligo dT Primer*. Next, this cDNA was used as the template in a PCR with the oligonucleotides *GeneRacer 5' Primer*, which represents the DNA equivalent of the 5' end of the *GeneRacer RNA Oligo*, and *LUC-4* which hybridizes to the coding-strand of the luciferase gene. The resulting PCR products were then used as templates in a second round of PCR using the oligonucleotides *GeneRacer 5' Nested Primer*, which represents the DNA equivalent of the 3' end of the *GeneRacer RNA Oligo*, and *LUC-2* which hybridizes to the coding-strand of the luciferase gene 5' to the *LUC-4* hybridization site. The products of this second reaction were then cloned using the TOPO TA Cloning Kit for Sequencing (Invitrogen Life Technologies, Carlsbad, CA). Clones containing the luciferase coding region were identified by filter hybridization. The radiolabeled probe used in this analysis was a 720 bp *EcoRI/PstI* fragment isolated from the plasmid p43Wt/Zeo which spans the CD43 promoter and the 5' end of the luciferase reporter gene. Clones containing the luciferase coding region were analyzed by DNA sequencing. The nucleotide sequences of the oligonucleotides used in RT-PCR were:

- 53 -

GeneRacer RNA Oligo:

5'-CGACUGGAGCACGAGGACACUGACAUGGACUGAAGGAGUAGAAA-3' (SEQ ID NO: 24)

GeneRacer Oligo dT Primer:

5'-GCTGTCAACGATACGCTACGTAACGGCATGACAGTG(T)₁₈-3'. (SEQ ID NO: 25)

GeneRacer 5' Primer:

5'-CGACTGGAGCACGAGGACACTGA-3' (SEQ ID NO: 26)

GeneRacer 5' Nested Primer:

5'-GGACACTGACATGGACTGAAGGAGTA-3' (SEQ ID NO: 27)

LUC-2:

5'-ATAGCCTTATGCAGTTGCTCT-3' (SEQ ID NO: 23)

LUC-4 :

5'-CACTACGGTAGGCTGCGAAATGTTCACTGTT-3'. (SEQ ID NO: 28)

15 *Transfection:* K562 cells were transfected by electroporation as previously described (Potter *et al.*, (1984) *Proceedings of the National Academy of Sciences USA*, 81:7161-7165.

59. Shelley *et al.*, (1991) *Proceedings of the National Academy of Sciences USA*, 88:10525-10529). Cells were transiently transfected with 23 µg of p43 Wt together with 2

20 µg of the plasmid pRSV-β (Promega Corp., Madison, WI) which contains the *lacZ* gene encoding β-galactosidase. Each transfection of p43 Wt was performed in parallel with a transfection of the promoterless luciferase plasmid pATLuc. Transfected cells were then either left untreated or treated with PMA for 12 hours before harvesting and lysis.

Luciferase and β -galactosidase activities were subsequently determined using reagents

25 purchased from Promega. Corp. (Madison, WI) and Tropix Inc. (Bedford, MA), respectively. Luciferase and β-galactosidase activities, assessed as light output, were measured using a Moonlight Luminometer which integrated peak luminescence 10

seconds after injection of assay buffer. The levels of β-galactosidase activity resulting from different transfections were taken as reflective of relative transfection efficiency and

30 used to correct the measurements of luciferase activity. Trans-repression by hnRNP-K in PMA treated K562 cells was assessed by transient transfections in which 8 µg of pATLuc

- 54 -

or p43Wt were mixed with 1 µg of pRSV-β and 16 µg of either Full-Length hnRNP-K or the equivalent vector empty of hnRNP-K coding sequences. Full-Length hnRNP-K contains the human hnRNP-K coding region downstream of the cytomegalovirus (CMV) promoter. After correction for transfection efficiency, the level of luciferase activity directed by pATLuc in the presence of Full-Length hnRNP-K was used to divide the levels of luciferase activity directed by p43Wt also in the presence of Full-Length hnRNP-K. This calculation yielded the fold above background activity of p43Wt in the presence of hnRNP-K. Equivalent calculations of luciferase activity in the presence of the CMV vector empty of the hnRNP-K coding region assessed non-specific effects caused by the vector backbone. The results (figures 5 and 6) from this second set of calculations represented the fold above background activity of p43Wt in the absence of hnRNP-K. K562 cells carrying stably within their genome the CD43 promoter linked to the luciferase reporter were transfected with 2 µg of pRSV-β and 23 µg of either Full-Length hnRNP-K or the equivalent vector empty of hnRNP-K coding sequences.

Affinity Purification: 200 µg of Streptavidin MagneSphere paramagnetic particles (Promega, Madison, WI) were washed two times in 1 ml of phosphate buffered saline (PBS) containing 0.1% bovine serum albumin (BSA) then mixed on a rotating wheel for 30 min with 1 ml of 0.2 M NaCl containing 10 µg of a version of the oligonucleotide CD43 PyRo SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267) which was biotinylated at its 3' end. The particles were then washed twice in 1X washing buffer (75 mM NaCl, 20 mM Tris-HCl pH 7.5, 1 mM EDTA, 15% glycerol, 0.05% IGEPAL CA-630). DNA affinity particles were mixed for 30 min at room temperature on a rotating wheel with 500 µl of binding mixture consisting of 1X binding buffer (70 mM KCl, 5 mM NaCl, 20 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 mM DTT and 10% glycerol), 300 µg of nuclear extract prepared from Jurkat cells and 0.2 µg/µl poly(dI-dC). The particles were captured with a magnetic stand and washed three times with 1 ml of 2X binding buffer containing 0.2 µg/µl poly(dI-dC). Bound protein was then eluted by adding 50 µl of washing buffer containing 1 M NaCl. The eluted protein was then subjected to EMSA and SDS-PAGE analysis.

Protein Identification by MALDI-TOF-MS: Affinity purified protein from Jurkat nuclear extracts was subjected to SDS-PAGE and stained with Coomassie blue. The

- 55 -

stained band was excised from the gel and then cut into small but uniform pieces. The gel was dehydrated with acetonitrile and rehydrated with 100 mM ammonium bicarbonate. Protein was protected from oxidation by incubation at 56°C for 1 hour with 10 mM DTT and the amino terminus protected by treatment with 10 mM iodoacetamide in 100 mM ammonium bicarbonate. Next, gel pieces were subjected to two rounds of washing with ammonium bicarbonate and subsequent drying with acetonitrile followed by a 12 hour incubation at 37°C with 12.5 ng/μl of trypsin in 50 mM ammonium bicarbonate (Rosenfeld *et al.*, (1992) *Analytical Biochemistry*, 203:173-179; Wilm *et al.*, (1996) *Analytical Chemistry*, 68:1-8). The masses of the trypsin-digested peptides were determined by matrix assisted laser desorption/ionization time-of-flight mass spectroscopy (MALDI-TOF-MS) using a Voyager DE-PRO instrument (Perceptive Biosystems Inc., Framingham, MA). Protein identification was achieved by- mass fingerprinting using the Mascot database created by Matrix Science.

Expression and Purification of Recombinant GST Proteins: The construct GST-RNP-K and its parent vector empty of hnRNP-K sequences were introduced into the *E. coli* strain XL-2 (Stratagene, La Jolla, CA). Bacteria were grown in 50 mls of Lauria broth at 37°C until they reached an optical density at 600 nm of 0.6 then recombinant protein expression was induced for 7 hours with 0.5 mM isopropyl β-D-thiogalactopyranoside (IPTG). Bacteria were washed twice in PBS and resuspended in 1 ml of 1X binding buffer (70 mM KCl, 5 mM NaCl, 20 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 mM DTT and 10% glycerol) containing "Complete" proteinase inhibitors (Roche Diagnostics Corp., Indianapolis, IN). Bacterial lysates were prepared by sonication and clarified by centrifugation. These preparations were then incubated for 30 min at room temperature with 200 μl of Glutathione-Sepharose beads (Amersham Pharmacia Biotech, Piscataway, NJ). Beads were washed three times with, 1 ml of PBS containing "Complete" inhibitors and bound proteins eluted by adding 800 μl of 50 mM Tris-HO, pH 8.0, 5 mM reduced glutathione. Finally, proteins were concentrated by centrifugation on a centricon-30 filter unit (Millipore Corp., Bedford, MA) for 15 min at 5000g.

Electrophoretic Mobility Shift Assay (EMSA): Oligonucleotides were labeled using T4 polynucleotide kinase and purified through Micro Bio-Spin 6 columns (Bio-Rad

- 56 -

Laboratories, Hercules, CA). DNA-protein binding reactions were carried out in a 20 μ l volume. Protein preparations were incubated with or without a molar excess of unlabeled specific or non-specific competitor oligonucleotides at 4°C for 10 min in 1X binding buffer (70 mM KCl, 5 mM NaCl, 20 mM Tris-HCl pH 7.5, 0.5 mM EDTA, 1 mM DTT and 10% glycerol) with 2.4 μ g/ μ l poly(dI-dC). Radiolabeled oligonucleotides were then added and the incubation continued for 20 min. DNA-protein complexes were resolved by electrophoresis through 4 or 6% polyacrylamide gels with 0.5X TBE and visualized by autoradiography. The oligonucleotides used in the EMSA analyses were:

CD43 PyRo SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267):

5'-GGGCCCCTTCCTTTCCCCTTG-3' (SEQ ID NO: 19)

CD43 PyRo SSUB: 5'-GGGCCCACUCCCUUCCCCUUGB-3' (SEQ ID NO: 20)

(U = Bromouracil; B = Biotin).

CD43 Mut-11 (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267):

5'-GGGCCCCTTCCTTCATATATG-3' (SEQ ID NO: 21)

NS-SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267):

5'-GAGTTAGCTCACTCATTAGG-3' (SEQ ID NO: 22)

UV Cross-Linking: UV crosslinking was performed using the oligonucleotide CD43 PyRo SSUB. This oligonucleotide is the same as the PyRo1 binding probe CD43 PyRo SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267) except that it is biotinylated at its 3' end and its deoxythymidines have been substituted with 5-bromo-2'-deoxyuridine. 30 μ g of Jurkat nuclear extract were preincubated for 5 minutes in 50 μ l of EMSA binding buffer containing 2.4 μ g/ μ l poly(dI-dC) then 10⁵ cpm of radiolabeled CD43 PyRo SSUB were added and binding performed for 20 minutes on ice in 1.5 ml microcentrifuge tubes. The tubes were then opened and placed in a UV crosslinker (Stratagene, La Jolla, CA) and exposed to 254 nm UV light for 10 minutes. Next 100 μ l of washed streptavidin-coated magnetic particles were added and crosslinked CD43 PyRo SSUB-protein complexes captured by rotation for 20 minutes at room temperature. Streptavidin particles were sedimented with a magnetic stand, washed five times with 500 μ l of EMSA binding buffer and then resuspended in 25 μ l of EMSA buffer containing 2% SDS and 5% 2-mercaptoethanol. Samples were boiled for 5 minutes,

- 57 -

centrifuged to pellet the magnetic particles and the supernatant analyzed by electrophoresis through a 12% SDS-polyacrylamide gel and subsequent autoradiography.

Results

Down-Regulation of CD43 mRNA Levels During K562 Activation: During
5 activation leukocytes increased their adhesive capacity mediated in part by a
down-regulation of CD43 expression. We have shown that 48 hours after activation of the
monocytic cell line U937 with phorbol ester there was a 69% reduction in CD43 mRNA
levels (Shelley *et al.*, (2001) *Br Journal of Haematology*, 115:159-166). A more dramatic
reduction in CD43 mRNA levels followed activation of the pre-erythroid/pre-
10 megakaryocytic cell line K562 (Figure 1).

Total RNA was prepared from K562 cells treated with PMA for the times
indicated. RNA was then subjected to northern blot analysis using an *EcoRI/NcoI*
fragment (CEM-E/N) isolated from the CD43 cDNA clone pCEM1.7 which has previously
been used to detect CD43 mRNA (Shelley *et al.*, (1989) *Proceedings of the National*
15 *Academy of Sciences USA*, 86:2819-2823). As a control for RNA loading, the same
northern blot hybridized with the CD43 specific probe was subsequently hybridized with a
probe which specifically recognizes GAPDH mRNA (Tso *et al.*, (1985) *Nucleic Acids*
Research, 13:2485-2502). Within 6 hours of K562 cells being activated with phorbol ester
the steady-state levels of CD43 mRNA become barely detectable.

Repression of CD43 Promoter Activity During K562 Activation: Previous studies
20 have indicated that the *cis*-acting elements responsible for directing the leukocytic
expression of the CD43 gene may be located in the proximal promoter region (Shelley *et*
al., (2001) *Br Journal of Haematology*, 115:159-166; Farokhzad *et al.*, (2000) *Nucleic*
Acids Research, 28:2256-2267; Kudo *et al.*, (1991) *Journal of Biological Chemistry*,
25 266:8483-8489; Kudo *et al.*, (1994) *European Journal of Biochemistry*, 223:319-327). The
construct p43Wt containing nucleotides -2 to +99 of the CD43 gene promoter was
transfected into K562 cells along with the control plasmid pRSV- β encoding β -
galactosidase. Transfected cells were then either left untreated (-PMA) or treated for 12
hours with PMA (+PMA) prior to harvesting. Luciferase values were measured and
30 normalized against β -galactosidase levels to correct for transfection efficiency.

- 58 -

The transfection experiments (employing the expression construct p43Wt which contains the CD43 promoter spanning nucleotides -2 to +99) indicated that the proximal promoter also contained the elements required for CD43 repression during K562 activation. Specifically, we found that expression of p43Wt was repressed by 39% upon PMA mediated activation of K562 cells (Figure 2). Expressed as histograms are the levels of luciferase gene activity directed by p43Wt in untreated and PMA treated K562 cells after division by the background activity conferred by the control plasmid pATLuc. Each histogram represents the mean \pm the standard deviation of three independent transfection experiments.

Identification of a Putative Repressor of the CD43 Promoter: Initially we have set out to determine the relationship between PyRol and hnRNP-K. We have discovered herein that PyRol and hnRNP-K are likely the same molecule (or at least structurally similar) and that these molecules are useful for repressing the CD43 promoter. In order to determine the function of PyRol we undertook its molecular cloning. PyRol was purified from Jurkat T-lymphocytic cells by affinity capture, digested with trypsin and its resulting peptides subjected to analysis by mass spectrometry. The fragmentation patterns produced from the PyRol peptides were then compared to those of known proteins deposited in the Mascot database created by Matrix Science. The most significant match to a known human protein was with heterogeneous nuclear ribonucleoprotein K (hnRNP-K) (Matunis *et al.*, (1992) *Molecular and Cellular Biology*, 12:164-171). Next, a fusion protein of hnRNP-K and glutathione S-transferase (GST) was produced in bacteria.

Figure 3 shows a radiolabeled single-stranded oligonucleotide, CD43 PyRo SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267), representing nucleotides +18 to +39 of the sense strand of the CD43 gene was incubated with no protein (Probe), purified GST (GST) or purified GST/hnRNP-K fusion protein (GST-K). Binding reactions containing GST/hnRNP-K were performed in the absence (GST-K) or presence (+) of a 100 fold molar excess of unlabeled CD43 PyRo SS, the presence of Mut-11 (Mut) representing a mutant version of CD43 PyRo SS which fails to support PyRol binding Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267) or the presence of an unrelated oligonucleotide (NS). The free probe and probe bound by the GST/hnRNP-K protein (GST:hnRNP-K) are arrowed.

The fusion protein was purified and shown to bind a radiolabeled single-stranded oligonucleotide representing the PyRol binding site within the CD43 gene promoter (Figure 3). This binding was effectively competed by an unlabeled excess of the PyRol binding site. However, binding failed to be competed with an identical molar excess of a mutant version of the binding site which previously failed to support PyRol interaction (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267). These studies demonstrate that the DNA binding characteristics of PyRol and hnRNP-K are similar. In addition, UV crosslinking demonstrated that the apparent molecular mass of PyRol is in the range of 50-65 kDa (Figure 4) which is comparable to that reported for hnRNP-K (Dejgaard *et al.*, (1994) *Journal of Biological Chemistry*, 236:33-48).

The left panel in Figure 4 shows the EMSA analysis performed as described in Fig. 3 except binding reactions contained the radiolabeled oligonucleotide CD43 PyRo SSUB which is both bromouracil and biotin modified and either no protein extract (Probe) or a nuclear extract prepared from Jurkat cells. Binding reactions containing Jurkat nuclear extract were performed in the absence (-) or presence (+) of a 100 fold molar excess of unlabeled CD43 PyRo SS (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267) or the presence of an unrelated oligonucleotide (NS). Marked with arrows are the positions of migration through a native polyacrylamide gel of CD43 PyRo SSUB unbound by protein (Free Probe) and bound by hnRNP-K. This analysis established that the protein binding characteristics of CD43 PyRo SSUB are indistinguishable from its equivalent CD43 PyRo SS which is modified neither with bromouracil nor biotin (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267) (right panel of Figure 4). The oligonucleotide CD43 PyRo SSUB was radiolabeled, incubated with (-) or without (Probe) a nuclear extract prepared from Jurkat cells and then exposed to UV light. CD43 PyRo SSUB was captured by streptavidin-coated magnetic particles, washed and subjected to electrophoresis through a 12% SDS-polyacrylamide gel and autoradiography.

Originally, hnRNP-K was identified as a RNA binding protein and recently its consensus RNA binding site was determined as UC₃₋₄(U/A)₂ (Matunis *et al.*, (1992) *Molecular and Cellular Biology*, 12:164-171; Thisted *et al.*, (2001) *Journal of Biological Chemistry*, 276:17484-17496). The single-stranded DNA equivalent of this consensus is present in the PyRol binding site within the CD43 promoter and mutation analysis has

- 60 -

demonstrated that this sequence is critical for PyRol binding (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267). Consequently, by the criteria of molecular cloning, molecular mass analysis and nucleic acid binding characteristics PyRol and hnRNP-K are structurally similar.

5 *Repression of the CD43 Gene Promoter by hnRNP-K.* We sought to determine if hnRNP-K could indeed function in such a capacity. Transfection of 16 µg of a plasmid expressing hnRNP-K indicated that in PMA treated K562 cells hnRNP-K did indeed represses the transcriptional activity of p43Wt. The level of this repression averaged 38% (Figure 5).

10 Eight micrograms of the luciferase reporter construct p43Wt were transfected into K562 cells mixed with either 16 µg of Full-Length hnRNP-K which expresses hnRNP-K or 16 µg of its parent vector empty of hnRNP-K coding sequences. One microgram of the β-galactosidase expression plasmid pRSV-β was also included in each transfection to control for transfection efficiency. Transfected cells were treated with PMA for 12 hours,
15 harvested and luciferase and galactosidase assays performed. The levels of β-galactosidase activity were taken as reflective of transfection efficiency and used to correct the luciferase assay results. Depicted as histograms are the levels of luciferase activity directed by p43Wt divided by those directed by the empty vector pATLuc in the presence of Full-Length hnRNP-K (+hnRNP-K) or the equivalent empty vector (-hnRNP-K). The
20 means of these levels ± the standard deviation resulting from three independent experiments are displayed in Figure 5.

It has been reported that the affects of hnRNP-K on transcriptional activity are dependent upon chromosomal structures (Lau *et al.*, (2000) *Journal of Cellular Biochemistry*, 79:395-406). Consequently, we generated a pool of K562 cell lines in
25 which the -2/+99 CD43 promoter linked to the luciferase reporter was randomly integrated into the genome.

K562 cells were transfected with the linearized plasmid p43Wt/Zeo, which was derived from p43Wt by insertion of the zeocin resistance gene. K562 cells in which the p43Wt/Zeo plasmid was stably integrated within the genome were selected by treatment
30 with zeocin. The mixed pool of zeocin resistant cells were then transfected with 2µg of pRSV-β mixed with either 23 µg of Full-Length hnRNP-K or 23 µg of its parent vector

- 61 -

empty of hnRNP-K coding sequences. Transfections were also performed using 9 µg of pRSV-β mixed with either 16 µg of Full-Length hnRNP-K or 16 µg of its parent. In those experiments in which the role of DNA methylation was assessed, cells were either untreated or pretreated for 48 hours with 5-azacytidine prior to transfection with 2 µg of pRSV-β mixed with either 23 µg of Full-Length hnRNP-K or 23 µg of its parent. Following all transfections cells were treated with PMA for 12 hours prior to harvesting and assay of luciferase and β-galactosidase activities. In experiments assessing the influence of DNA methylation, transfected cells, untreated or pretreated with 5-azacytidine, were treated simultaneously with PMA and 5-azacytidine for 12 hours prior to harvesting. The levels of β-galactosidase activity were taken as reflective of transfection efficiency and used to correct the luciferase assay results. The level of luciferase activity directed by the CD43 promoter in the presence of the vector empty of HnRNP-K coding sequences (-hnRNP-K) was assigned a value of 100%. The luciferase levels directed by the CD43 promoter matched, parallel transfections employing Full-Length hnRNP-K (+hnRNP-K) was calculated as a proportion of the 100% value. The mean of these proportional values ± the standard deviation resulting from three independent experiments are displayed in Figure 6.

Transfection of the hnRNP-K expression plasmid into this stable cell line subsequently treated with PMA repressed luciferase reporter gene activity by an average of 62% (Figure 6). Consequently, hnRNP-K significantly represses the CD43 promoter either in a chromosomal or extra chromosomal context.

Repression Mediated by hnRNP-K is Dependent Upon DNA Methylation. Previous studies have demonstrated that in non-hematopoietic cells the CD43 promoter is maintained in an inactive state by DNA methylation (Kudo *et al.*, (1995) *Journal of Biological Chemistry*, 270:13298-13302; Kudo *et al.*, (1998) *Molecular and Cellular Biology*, 18:5492-5499). We tested whether hnRNP-K repression of the CD43 promoter was methylation dependent. The pool of stable cell lines containing the CD43 promoter linked to luciferase was treated for 48 hours with the DNA methyltransferase inhibitor 5-azacytidine (Creusot *et al.*, (1982) *Journal of Biological Chemistry*, 257:2041-2048). These treated cells were then transfected with the hnRNP-K expression plasmid and incubated for 12 hours in the presence of PMA and 5-azacytidine. Under these

- 62 -

circumstances the CD43 promoter was repressed by only 17% (Figure 6). Without a 48 hour pretreatment but with a 12 hour treatment with 5-azacytidine and PMA immediately after transfection hnRNP-K repressed the CD43 promoter by 50%. In cells treated only with PMA after transfection hnRNP-K represses the CD43 promoter by 62% (Figure 6).

- 5 These results indicate that exposure of the hnRNP-K expression plasmid to 5-azacytidine for 12 hours influences its ability to effect repression by only 12%. In contrast, exposure of the CD43 promoter to 5-azacytidine for 48 hours reduces the repressive capacity of hnRNP-K by 33%. Consequently, repression of the CD43 promoter by hnRNP-K is at least partially affected by this promoter being methylated.

- 10 *Determination of the Transcription Start Site of the CD43 Gene when Linked to Luciferase and Integrated within the Genome of K562.* Transcription of the CD43 gene has been shown to be initiated at two major sites (Shelley *et al.*, (1990) *Biochemical Journal*, 270:569-576) between which lies the binding site for hnRNP-K. hnRNP is known to bind RNA and has been implicated in mediating RNA stability and transport (Pi-ol-Roma *et al.*,
15 (1992) *Nature*, 355:730-732; Pi-ol-Roma *et al.*, (1993) *Trends in Cell Biology*, 3:151-155; Michael *et al.*, (1995) *Cold Spring Harbor Symp Quant Biology*, 60:663-668; Kiledjian *et al.*, (1995) *EMBO Journal*, 14:4357-4364; Visa *et al.*, (1996) *Cell*, 84:253-264; Michael *et al.*, (1997) *EMBO Journal*, 16:3587-3598; Dreyfuss *et al.*, (2002) *Nature Reviews: Molecular Cell Biology*, 3:195-205). Consequently, if transcription of the CD43/luciferase
20 gene is initiated at the upstream of the two potential start sites, the affect of hnRNP-K on luciferase activity may reflect mechanisms mediated by RNA binding rather than transcriptional control mechanisms. Due to this reasoning, we identified the transcription start site of the CD43/luciferase gene present in the genome of K562 using RT-PCR coupled with RACE technology. Southern blot analysis of the RT-PCR products resulting
25 from this procedure revealed a single discrete cDNA containing the luciferase coding region (Figure 7). Cloning and sequencing of this cDNA determined that the downstream of the two potential transcription start sites is utilized by the CD43 promoter when linked to luciferase and integrated within K562. Therefore, the transcript produced from this promoter does not contain a binding site for hnRNP-K. Consequently, repression of the
30 CD43/luciferase fusion gene by hnRNP-K is likely mediated by transcriptional

- 63 -

mechanisms as opposed to mechanisms involving RNA binding such as decreased RNA stability and/or transport.

The mixed pool of K562 cells containing within their genome the CD43/luciferase fusion gene were treated for 24 hours with PMA and then total RNA isolated using an RNeasy Maxi Kit (Qiagen, Inc., Valencia, CA). A GeneRacer Kit (Invitrogen Life Technologies) was employed to ligate the *GeneRacer RNA Oligo* specifically to the 5' end of full-length mRNA which was then reverse-transcribed using the *GeneRacer Oligo dT Primer*. The resulting products were used as templates in a PCR containing the primer *LUC-4*, which hybridizes to the coding strand of the luciferase gene, and the *GeneRacer 5' Primer*, which represents the DNA equivalent of the 5' end of the *GeneRacer RNA Oligo*. PCR products were then subjected to Southern blot analysis using as probe the radiolabeled oligonucleotide *LUC-2* which hybridizes to the coding strand of the luciferase gene further 5' than does *LUC-4*. The hybridization signal detected by autoradiography is depicted in Figure 7

HnRNP-K Acts Together with Pura to Effect Repression of the CD43 Promoter.

We have previously demonstrated that *Pura* interacts with the CD43 promoter immediately upstream of the region which we show here binds hnRNP-K (Shelley *et al.*, (2001) *Br Journal of Haematology*, 115:159-166). In the pro-monocytic cell line U937 treated with PMA *Pura* represses the CD43 promoter. Therefore, it was of interest to determine whether *Pura* also mediates repression within K562 cells and whether it works with or against hnRNP-K. In transfection experiments employing expression plasmids constitutively expressing hnRNP-K or *Pura* we found that alone *Pura* represses the CD43/luciferase gene present within the K562 genome by 36% (Figure 8). However, when recombinant *Pura* is expressed together with recombinant hnRNP-K repression of the CD43 promoter is increased to 51%. Consequently, hnRNP-K and *Pura* have the ability to work together to effect repression.

The β -galactosidase expression construct pRSV- β was mixed with one of four different combinations of plasmids. First, pHAPur1 and Full-Length hnRNP-K which express *Pura* and hnRNP-K, respectively. Second, pHAPur1 and the empty vector equivalent of Full-Length hnRNP-K. Third, Full-Length hnRNP-K and pHA which represents the empty vector equivalent of pHAPur1. Fourth, both empty vectors. These

- 64 -

four DNA mixtures were then transfected into the mixed pool of K562 cells containing the CD43/luciferase fusion and treated for 12 hours with PMA. Cells were harvested, lysed and luciferase and β -galactosidase assays performed. The levels of β -galactosidase activity were taken as reflective of transfection efficiency and used to correct the luciferase assay results. Using these corrected values, the level of luciferase activity directed by the CD43 promoter in the presence of the empty vectors was assigned a value of 100%. The levels of luciferase activity directed by the promoter in the presence of constructs expressing Puro α and/or hnRNP-K was calculated as a proportion of the 100% value. The mean of these levels + the standard deviation resulting from three independent experiments are displayed as histograms (Figure 8).

On the surface of resting leukocytes CD43 prevents cellular interaction. However, during leukocyte activation the glycosylation pattern of CD43 changes and there is a down-regulation of the overall amount of CD43 expressed on the cell surface. These alterations in both the qualitative and quantitative expression of CD43 result in the promotion of both homotypic and heterotypic leukocyte interactions. Using *in vitro* models of leukocyte activation we have found that down-regulation of CD43 is mediated by repression of the transcriptional activity of the CD43 gene promoter. Previously, we demonstrated that the transcription factor Puro α mediates repression of the CD43 promoter (Shelley *et al.*, (2001) *Br Journal of Haematology*, 115:159-166). We have found according to some aspects of the invention that hnRNP-K also mediates CD43 gene repression. The binding sites for Puro α and hnRNP-K lie adjacent to one another within the CD43 promoter. Puro α interacts with the -2/+17 region of the CD43 gene and hnRNP-K interacts with the +18/+39 region. Both factors are induced during U937 activation concomitant with repression of CD43 promoter activity.

Puro α and hnRNP-K interacted preferentially with single-stranded DNA indicating that single-stranded structures within the CD43 promoter are probably important in regulating its activity. Nuclease S1 analysis has established that single-stranded structures can exist within the CD43 promoter (Farokhzad *et al.*, (2000) *Nucleic Acids Research*, 28:2256-2267). Furthermore, analysis of the primary sequence of the promoter revealed repeats which could form *slippage* or *cruciform* structures (Yagil *et al.*, (1991) *Critical Review of Biochemistry Molecular Biology*, 26:475-559). The single-stranded loop of a

- 65 -

slippage structure within the -2/+17 region would support efficient Pur α binding while the loop of a *cruciform* structure within the +18/+39 region would support efficient binding of hnRNP-K (Figure 9). The hnRNP-K protein was originally characterized as an RNA binding protein.

5 Figure 9 shows the misalignment of direct repeats of the sequence GGTGG may form *slippage* structures at the 5' end of the promoter. The single-stranded region on the sense-strand of these structures would support efficient binding of Pur α . Two types of *slippage* structures are possible. One is illustrated. Pairing of an inverted repeat of the sequence CAGGGCCC could form a *cruciform* structure immediately downstream of the

10 *slippage* structures. This possible *cruciform* is illustrated. The single-stranded loop region on the sense strand of the *cruciform* would support efficient binding of hnRNP-K. The DNA equivalent of the consensus RNA binding site of hnRNP-K is marked by bold characters (Thisted *et al.*, (2001) *Journal of Biological Chemistry*, 276:17484-17496). The nucleotides defined by mutation analysis as being important for hnRNP-K binding are

15 marked with arrows. The two major transcription initiation sites of the CD43 gene are marked with asterisks.

The site within the CD43 promoter which binds hnRNP-K lies adjacent to a site which interacts with Sp1 (Shelley *et al.*, (2001) *Br Journal of Haematology*, 115:159-166; Kudo *et al.*, (1994) *European Journal of Biochemistry*, 223:319-327; Kudo *et al.*, (1998)

20 *Molecular and Cellular Biology*, 18:5492-5499). This Sp1 binding site overlaps the region which binds Pur α . Transient transfections have indicated that, unlike hnRNP-K and Pur α , Sp1 functions to activate the CD43 promoter 63. Since Sp1 binds double-stranded DNA a mechanism of repression can be envisioned in which a double-stranded CD43 promoter supporting Sp1 binding and transcriptional activation acquires a single-stranded

25 conformation supporting hnRNP-K and Pur α binding. Such opening of the CD43 promoter may in fact be mediated by Pur α and hnRNP-K themselves. This possibility was supported by the finding that Pur α is capable of displacing an oligonucleotide annealed to single-stranded M13 DNA (Darbinian *et al.*, (2001) *Journal of Cellular Biochemistry*, 80:589-595). In addition, hnRNP-K has been shown to impart a single-stranded

30 conformation to the CT element of the *c-myc* promoter contained within a supercoiled plasmid (Michelotti *et al.*, (1996) *Molecular and Cellular Biology*, 16:2656-2669).

- 66 -

However, in the context of the *c-myc* promoter hnRNP-K acted as a transcriptional activator not a repressor as it did within the CD43 promoter (Michelotti *et al.*, (1996) *Molecular and Cellular Biology*, 16:2350-2360; Takimoto *et al.*, (1993) *Journal of Biological Chemistry*, 268:18249-18258). Therefore, while DNA unwinding may be
5 intrinsic properties of Pur α and hnRNP-K, the functional consequences of this activity appeared dependent upon promoter architecture and probably also cellular environment.

Previous studies have shown that in non-hematopoietic cells the CD43 gene is maintained in a transcriptionally inactive state by methylation of CpG dinucleotides within the promoter (Kudo *et al.*, (1995) *Journal of Biological Chemistry*, 270:13298-13302).

10 These methylated dinucleotides are then able to bind the methyl-CpG binding protein MeCP2 (Kudo *et al.*, (1998) *Molecular and Cellular Biology*, 18:5492-5499). We have shown that repression of the CD43 promoter by hnRNP-K may be affected by DNA methylation. This finding suggests that repression of the CD43 promoter during hematopoietic differentiation may be mediated by an increase in its degree of methylation.
15 Such an increase could result in the recruitment of methyl-CpG binding proteins and/or the more effective recruitment of hnRNP-K.

Example 2

Introduction

20 Ovarian cancer is the leading cause of gynecological cancer deaths and the fifth leading cause of all cancer deaths among American women. An encouraging trend of slightly decreasing mortality rates has been due largely to advances in early detection. In contrast, we have failed to significantly advance treatment beyond the *sledge-hammer* approaches of surgery, chemotherapy and radiotherapy.

25 Meaningful advances in ovarian cancer treatment will stem only from identifying new diagnostic and prognostic tools and targeting the causative molecular defects.

Recently, we have identified abnormal expression of the sialoglycoprotein CD43 as a novel defect characterizing ovarian cancer tissue. Normal ovarian cells fail to exhibit CD43 expression and no CD43 expression was observed on cells immediately next to the
30 malignant lesions.

- 67 -

CD43 is a large, abundant and highly charged trans-membrane molecule that is normally only expressed on the surface of white blood cells. In this normal context of expression CD43 acts as a barrier molecule preventing intercellular interactions and allowing blood cells to remain in the circulation.

5 *In vitro* evidence indicates that, as on white blood cells, CD43 also serves a barrier function on the surface of ovarian cancer cells. In this abnormal context of expression CD43 may prevent interaction with immune effector cells such as natural killers and, therefore, facilitates an escape from immunosurveillance.

As well as allowing malignant cells to avoid immune recognition, expression of
10 CD43 may also contribute to the pathogenesis of ovarian cancer in a another way. By inhibiting cell-cell interactions, CD43 could aid in turning the primary tumor into a loose cellular mass which sheds potentially metastatic neoplastic cells into the circulation.

Demonstration that CD43 is Abnormally Expressed in Ovarian Cancer Cells.

Human CD43 is a large, abundant and highly charged transmembrane molecule which
15 inhibits intercellular interaction (Remold-O'Donnell, *et al.*, (1990) *Immunodeficiency Rev.*, 2: 151-174; Ardman *et al.*, (1992) *Proc. Natl. Acad. Sci USA*, 89: 5001-5005; Manjunath *et al.*, (1993) *J. Immunol.*, 151: 1528-1534). Normally CD43 expression is restricted to the surface of hematopoietic cells. However, we have found that primary ovarian cancer tumors also exhibit CD43 expression. Snap frozen ovarian cancer tissue was obtained
20 from the Massachusetts General Hospital Tumor Bank, sectioned and stained using the monoclonal antibody BS1 which specifically recognizes CD43 independent of its glycosylation status and as secondary antibody donkey anti-Mouse Cy3.. The eight specimens analyzed consisted of two endometrial carcinomas, two clear cell carcinomas, two histologically undefined adenocarcinomas and two areas of patient-matched normal
25 ovarian tissue. All the carcinomas stained positive with the BS1 antibody(Figure 10 - panel #1 CA: endometrial ovarian carcinoma; panel #2 CA: clear cell ovarian carcinoma). However, staining of the undefined adenocarcinomas was particularly striking(Figure 10 - panel #3 CA: undefined ovarian adenocarcinoma). We detected no staining of the patient-matched normal ovarian tissue nor of the tissue surrounding the tumors (Figure 10 -
30 left panel: normal ovarian tissue from an ovarian cancer patient).

- 68 -

Demonstration that CD43 Expression Protects Cancer Cells From Natural Killing.

In 1995 it was demonstrated by the laboratory of Dr. Gunnar Hansson in Gothenburg, Sweden that CD43 is produced by the colon adenocarcinoma cell line COLO205 (Baeckstrom, *et al.*, (1995) *I. Rial. Chein.*, 270: 13688-13692). This was the first
5 identification of a CD43 molecule expressed outside the hematopoietic cell lineage. In 1996 the association of carcinoma with abnormal CD43 expression was confirmed by the laboratory of Dr. Ignacio Molina in Granada, Spain (Santamaria *et al.*, (1996) *Cancer Res.*, 56: 3526-3529). In 1997 the Hansson laboratory published a study which described the functional significance of CD43 expression by carcinoma cells. Returning to their
10 COLO205 model, the Swedish group isolated the CD43 produced by these cells and demonstrated it was able to specifically inhibit cytotoxic lysis mediated by natural killer cells (Zhang *et al.*, (1997) *Cell. Immunol.*, 176: 158-165). Inhibition was concentration dependent and at the highest concentration used (100 µg/ml) reached approximately 60%. This finding indicates that CD43 expression likely contributes to carcinogenesis by helping
15 malignant cells evade immune recognition.

Evasion of immunosurveillance by cancer cells is probably caused by CD43 providing an anti-adhesion shield of protection. Such an anti-adhesion function of CD43 is supported by three sets of experiments. In the first, introduction of CD43 into CD43-negative cervical carcinoma cells was shown to reduce CD54 mediated adhesion (Ardman
20 *et al.*, (1992) *Proc. Natl. Acad. Sci USA*, 89: 5001-5005). In the second, targeted disruption of the CD43 gene in cells derived from acute T-cell leukemia was shown to enhance homotypic adhesion and binding to fibronectin and mv-1 gpl20 (Manjunath *et al.*, (1993) *I. Immunol.*, 151: 1528-1534). Finally, in a third study, transgenic mice were produced in which the CD43 gene was specifically knocked-out. These mice exhibited
25 increased leukocyte adhesion and, as a consequence, premature clearance of leukocytes from the circulation (Dragone *et al.*, (1995) *Proc. Natl. Acad. Sci. USA*, 92: 626-630).

Taken together, work from around the world over the last few years has established the anti-adhesion function of CD43. It appears that cancer cells, by abnormally expressing CD43, may acquire this anti-adhesive function and, in so doing, acquire a means of
30 avoiding immunosurveillance.

- 69 -

Although not wishing to be bound to any particular theory or mechanism, we believe that abnormal expression of CD43 in ovarian cancer must be due either to mutation of the gene by which it is encoded and/or alterations in the transcription factors by which this gene is controlled. Our laboratory is in a unique position to identify these molecular defects since over the last decade we have both structurally and functionally characterized the CD43 gene.

Characterization of CD43 cDNA and Genomic Clones Initially we isolated the human CD43 molecule from the T-lymphoblastoid cell line CEM and determined the amino acid sequence of a 15 residue tryptic peptide. An oligonucleotide corresponding to the most frequently used codons specifying this peptide was then used to screen a CEM cDNA library. Sequencing of clones derived from this screen predicted that CD43 is an integral membrane protein with a 19 residue N-terminal hydrophobic leader peptide followed by a 235 residue extracellular region, a 43 residue transmembrane region and a 123 amino acid C-terminal intracellular region (Shelley *et al.*, (1989) *Proc. Natl. Acad. Sci. USA*, 86: 2819-2823). Southern blot and *in situ* hybridization analyses established that CD43 is encoded by a single gene located on the p 11.2 band of chromosome 16. This region is often rearranged in patients with acute myelomonocytic leukemia (Pallant *et al.*, (1989) *Proc. Natl. Acad. Sci. USA*, 86: 1328-1332; Hogge *et al.*, (1984) *I. Clin. Oncol.*, 2: 550-557; LeBeau *et al.*, (1983) *N. Engl. J. Med.*, 309: 630-636).

A human genomic library was screened with CD43 cDNA, a specific clone isolated and the nucleotide sequence of the entire CD43 gene determined (Shelley *et al.*, (1990) *Biochem. J.*, 270: 569-576). The gene is approximately 4.6kb in length and comprises two exons and one intron. S 1 nuclease protection and primer extension analysis demonstrated that transcription is initiated at two major sites. Upstream of the 5' major transcription initiation site, 990 nucleotides were sequenced. The promoter region was found to contain a number of short repeats but no *TATA* or *CAAT* elements.

The Proximal Promoter Region of the CD43 Gene Directs Tissue-Specific Expression. In order to determine the region of the CD43 gene responsible for driving appropriate cell-specific and tissue-specific expression, a series of deletion constructs were made. The smallest of these plasmid constructs was named p43Wt. This plasmid contains nucleotides - 2 to +99 of the CD43 gene fused to the 5' end of a promoterless luciferase

- 70 -

gene. The p43Wt construct was transfected into U937 pro-monocytic cells, Jurkat T-lymphocytes, K562 pre-erythroid/pre-megakaryocytic cells, CEM T-lymphocytic cells and the non-tumorigenic breast epithelial cell line MTSVI-7 (Bartek *et al.* (1991) *Proc. Natl. Acad. Sci. USA*, 88: 25320-2534). Measurement of luciferase activity demonstrated that in U937 cells p43Wt directs expression which is 338 fold higher than that directed by the parent plasmid. Such expression in U937 cells compares favorably to the 309 and 698 fold above parental expression levels directed by p43Wt in Jurkat T-lymphocytes and K562 pre-erythroid/pre-megakaryocytic cells, respectively. When p43Wt was transfected into CEM T-lymphocytic cells it directed a level of expression only 74 fold above that of its parent. This low level of promoter activity compared to other leukocytic cell types reflects the relatively low level of CD43 protein produced by CEM cells (Carlsson *et al.*, (1986) *J. Biol. Chem.*, 261: 12779-12786).

CD43 mRNA is undetectable in the epithelial cell line MTSV1-7. In this cell line p43Wt directs expression only 7 fold above that directed by the parental plasmid. Our findings differ from those of Kudo and Fukuda who reported the proximal CD43 promoter is active when transfected into epithelial cells (Kudo *et al.*, (1994) *Bur. J. Biochem.*, 223: 319-327). However, HeLa not MTSVI-7 cells were used in these studies. HeLa cells are derived from a cervical carcinoma (Gey *et al.*, (1952) *Cancer Res.*, 12: 264-265) while MTSVI-7 cells are derived from normal breast tissue (Bartek *et al.*, (1991) *Proc. Natl. Acad. Sci. USA*, 88: 3520-3524). The development of carcinoma is associated with dramatic changes in DNA methylation (Santini *et al.*, (2001) *Ann. Intern. Med.*, 134: 573-586; Schmutte *et al.*, (1998) *Biol. Chem.*, 379: 377-388). Kudo and Fukuda have reported that the tissue-specific expression of the CD43 gene is regulated by DNA methylation and its interaction with the methyl-CpG-binding protein MeCP2 (Kudo *et al.*, (1995) *J. Biol. Chem.*, 270: 13298-13302; Kudo *et al.*, (1998) *Mol. Cell. Biol.*, 18: 5492-5499). Therefore, it is possible that the inappropriate expression of the CD43 promoter when transfected into HeLa rather than MTSV1-7 cells reflects the disruption of the DNA methylation machinery associated with carcinogenesis.

In summary, we have shown that the region of the CD43 gene extending from -2 to +99 is sufficient *in vitro* to direct appropriate tissue-specific and cell-specific expression. This finding indicates the importance of analyzing the structure and function

- 71 -

of this region in ovarian cancer cells which exhibit inappropriate expression of the CD43 gene.

Example 3

5 Experimental evidence generated both *in vitro* and *in vivo* indicates that ovarian cancer is characterized by abnormal expression of CD43. Although not wishing to be bound to any particular theory or mechanism, we believe that abnormal expression of CD43 in ovarian cancer contributes in two major ways to disease progression. First, by inhibiting the engagement of ovarian cancer cells by immune effectors such as natural
10 killer cells, CD43 allows ovarian cancer cells to escape immunosurveillance. Second, by inhibiting adhesion, CD43 aids in turning the primary tumor into a loose cellular mass which sheds potentially metastatic neoplastic cells into the circulation. We believe that abnormal CD43 expression is of both diagnostic and prognostic value in ovarian cancer. We also believe that abnormal expression of CD43 in ovarian cancer would be due to
15 abnormal transcription of the gene by which it is encoded.

Determination of the Pattern of Abnormal CD43 Expression in Ovarian Cancer.

Our study of six ovarian cancer tumors has demonstrated that all six exhibit abnormal expression of CD43. This finding has led us to conclude that abnormal CD43 expression is of both diagnostic and prognostic value in ovarian cancer. A series of
20 immunocytochemical examinations using a panel of anti-CD43 antibodies are performed. Each of the distinct biological categories of ovarian tumors (benign, borderline and malignant), each of the histological subtypes (serous, mucinous, endometrioid and clear cell), each of the disease stages (Stage IA -Stage IV) and also BRCA1-associated hereditary ovarian cancer are examined. In addition, we examine a broad range of non-
25 tumorogenic and tumorogenic ovarian cells cultured *in vitro*. CD43 expression is evaluated in the primary and metastatic tumors which develop in each of seven newly established genetic mouse models of ovarian cancer.

Fluorescence microscopy is used to determine CD43 expression in three types of ovarian samples; sectioned human ovarian tumors, human ovarian cells cultured *in vitro*
30 and malignant cells generated within seven different genetic mouse models of ovarian cancer.

- 72 -

Human Ovarian Tumors: Fresh ovarian tumors and normal ovarian tissue are obtained from the *Gillette Center for Women's Cancers-Gynecologic Oncology* in collaboration with Arlan F. Fuller, Jr., M.D. and Robert L. DeBernardo, M.D. Snap frozen ovarian tumors and normal ovarian tissue are obtained from the *MGH Tumor Bank*. Using these sources of material we examine each of the distinct biological categories of ovarian tumors (benign, borderline and malignant), each of the major histological subtypes (serous, mucinous, endometrioid and clear cell) and also BRCA1-associated hereditary ovarian cancer. In addition, we will examine the different disease stages to determine whether CD43 expression is of prognostic value.

Human Ovarian Cells Cultured in Vitro: Three categories of cultured human ovarian cells are examined; non-tumorigenic primary cultures and cell lines, cystadenoma derived cell lines and ovarian carcinoma derived lines.

Non-tumorigenic cell lines:

- (1). Primary cultures established from normal ovarian epithelium (BioWhittaker, Inc., Walkersville, MD).
- (2). Cell lines derived by SV40-immortalization of normal ovarian epithelium; mO-106, mO-118, HIO-121, mo-135 IOSE-80 and IOSE-144 (Maines-Bandiera *et al.*, (1992) *Am. J. Obstet. Gynecol.*, 167: 729-725; Bruening *et al.*, (1999) *Cancer Res.*, 59: 4973-4983).
- (3). Cell lines derived by HPV16- immortalization of normal ovarian epithelium; HOSE-A and HOSE-B-(Gregoire *et al.*, (1998) *In Vitro Cell. Dev. Biol. Anim.*, 34: 636-639).

Cystadenoma-derived cell lines

- (1). ML3, MLI0, MCV39, MCV50 and MCV152 (Luo *et al.*, (1997) *Gynecol. Oncol.*, 67: 277-284).

Ovarian carcinoma-derived cell lines

- (1). Primary cultures of ovarian tumors established on a fee-for-service basis by BioWhittaker, Inc. from fresh material supplied by the *Gillette Center for Women's Cancers-Gynecologic Oncology*.
- (2). Caov-3, Caov-4, NIH OVCAR-3, SKOV3, ES-2, PA-1 and SW626 (American Type Culture Collection, Rockville, MD).

- 73 -

- (3). OVCAR-10, OVCAR-8, OVCAR-5, OVCAR-4, PEO1, PEO4, A1847 and A2780 (Godwin *et al.*, (1992) *Proc. Natl. Acad. Sci. USA*, 89: 3070-3074).
- (4). OVCA433, OVCA420, OVCA429, OVCA432 and DOV13 (Moser *et al.*, (1996) *Int. J. Cancer*, 67: 695-701).
- 5 (5). HEY, HOC-1, HOC-7 and HOC-8 (Buick *et al.*, (1985) *Cancer Res.*, 45: 3668-3676).
- (6). ML9A, ML12B and ML23A (Luo *et al.*, (1997) *Gynecol. Oncol.*, 67: 277-284).
- (7). HOSE-A and HOSE-B exhibiting spontaneous malignant transformation due to extended culture on collagen rafts (Gregoire *et al.*, (2001) *Clin. Cancer Res.*, 7: 4280-4287).
- 10

Genetic Mouse Models of Ovarian Cancer: Recently, mouse models of ovarian cancer have been developed in which avian retroviral vectors are used to introduce multiple genetic lesions into ovarian surface epithelial cells (Fisher, *et al.*, (1999) *Oncogene* 18: 5253-5260; Orsulic *et al.*, (2002) *Cancer Cell* 1: 53-62; Orsulic (2002) *Mamm. Genome*, 13: 543-547). To date, the expression of seven combinations of the oncogenes *c-myc*, *K-ras* and *akt* within p53-t- mice have been shown to produce ovarian tumors. The different combinations of oncogenes result in different frequencies of primary tumor development and in tumors with different metastatic potential. We examine CD43 expression on sectioned primary and metastatic tumors in each of the seven genetic models of ovarian cancer.

15

20

Staining Procedure: Sectioned human tissue and human cell lines growing on microscope slides are blocked with serum and then incubated with a mouse anti-human CD43 antibody or an isotype matched negative control antibody. A panel of anti-CD43 antibodies is used consisting of L10, L14, DFT1, BRA7G and BS1 which recognize the five different epitope clusters identified on the human CD43 molecule (Remold-O'Donnell (1995) *Leukocyte Typing V*, 1697-1701; De Smet *et al.*, (1995) *Leukocyte Typing V*, 1706-1709; Axelsson, *et al.*, (1995) *Leukocyte Typing V*, 1708-1709). Sectioned mouse tissue is analyzed using the goat anti-mouse CD43 antibody S19 (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) and the rat anti-mouse antibodies S7 and 1B11 (PharMingen, San Diego, CA). After washing in PBS the cells are incubated with FITC labeled anti-mouse immunoglobulin and observed using a fluorescence microscope. As controls for these

25

30

- 74 -

experiments immunocytochemical analysis is performed in parallel on CEM cells which express CD43 and on HeLa cells which do not. These studies determine diagnostic and prognostic value of CD43 expression in ovarian cancer.

Determination of CD43 Gene Structure in Ovarian Cancer. Large-scale gene rearrangements and amplifications which result in inappropriate gene expression have been shown to be associated with and, in some cases, to cause malignancy. The presence of gross structural and copy number defects of the CD43 gene in ovarian cancer is determined by Southern blot and/or Polymerase Chain Reaction (PCR) analysis. These analyses are performed on CD43-negative and CD43-positive ovarian cell cultures as well as on CD43-negative and CD43-positive ovarian tissue. On the tissue samples CD43 expression is visualized by immunocytochemistry and CD43-negative and CD43-positive cells isolated by laser capture microdissection (LCM). In Southern blot analysis, cell cultures or material isolated by LCM are embedded in agarose plugs and lysed in SDS. DNA is digested within the agarose plugs with combinations of the restriction enzymes *AscI*, *NotI*, *PacI*, *Sce83871* and *SfiI*. These enzymes have long recognition sequences that occur infrequently within the human genome and, therefore, produce large DNA fragments. Such fragments are resolved by Pulsed Field Gel Electrophoresis (PFGE) and transferred to nitrocellulose filters by Southern blotting (Southern, (1975) *J. Mol. Biol.*, 98: 503-517). After baking, the filter is hybridized with a ³²p-labeled CD43 cDNA probe, washed and subjected to autoradiography. The human genome database allows probes specific to the CD43 gene locus but distal to the CD43 coding region to be generated by the PCR technique. These probes are also employed in Southern blot analysis. In this way we produce a restriction map of the CD43 gene locus which spans hundreds of kilobases. The hybridization pattern resulting from CD43-positive and CD43-negative material is compared to evaluate chromosomal rearrangements. Quantitation of hybridization signals by densitometry evaluates chromosomal amplification. The large scale of the mapping possible using PFGE allows chromosomal rearrangements distal to the CD43 coding region to be detected. In order to avoid the possibility of detecting non-functional Restriction Fragment Length Polymorphisms hybridization patterns resulting from the digestion of a number of different restriction endonucleases are compared. Differences in patterns in CD43-positive compared to CD43-negative material judge and identify the

- 75 -

rearranged CD43 gene locus. This rearrangement is localized by performing Southern blot analysis using restriction enzymes such as *HindIII*, *BamHI* and *EcoRI* which have hexanucleotide recognition sequences.

In order to characterize any rearrangement of the CD43 gene detected by Southern blotting and also identify any small scale mutations invisible to this analysis, the CD43 gene is cloned from CD43- positive ovarian cancer cells and sequenced. First, total genomic DNA extracted from ovarian cells expressing CD43 is digested with the restriction enzymes that Southern blot analysis has determined produce fragments spanning the CD43 gene locus. These digests are subjected to agarose gel electrophoresis and the appropriately sized DNA isolated. The isolated fragments are then ligated into plasmid vectors and the CD43 clones, detected by hybridization screening, sequenced by standard means (Shelley *et al.*, (1989) *Proc. Natl. Acad. Sci. USA*, 86:2819-2823; Ausubel *et al.*, (1994) *John Wiley and Sons, Inc.*). The sequences of the CD43 genes isolated from CD43-positive ovarian cancer cells are compared with published sequences to identify any mutations. Sequences which cannot be assigned as being of CD43 gene origin due to gross chromosomal rearrangement are identified by BLAST searches of the Genbank nucleotide sequence database. These studies will identify abnormal expression of CD43 in ovarian cancer associated with mutation of the CD43 gene.

Small quantities of genomic DNA can be isolated by LCM. This limitation may preclude Southern blot analysis of CD43 gene structure in CD43-positive tumor tissue. Under these circumstances we evaluate the structure of the CD43 gene by linker modified (LM) PCR. Genomic DNA isolated by LCM is digested with a series of restriction enzymes which cut in or outside the transcribed region of the CD43 gene. Oligonucleotide linkers are then ligated to the cleavage sites and the resulting products are used as templates in PCR. One of the primers employed in these reactions hybridizes to the linker and the other to the known CD43 gene sequence on one side or the other of the restriction enzyme cleavage site. Comparison of the size of the PCR products generated from CD43-positive and CD43-negative tissue assesses rearrangement of the CD43 gene locus. Gene amplification is assessed by real-time quantitative PCR using TaqMan@ probes (Applied Biosystems, Foster City, CA).

- 76 -

CD43 gene expression has been shown to be dependent upon its pattern of methylation (Kudo *et al.*, (1995) *J. Bioi. Chem.*, 270: 13298-13302; Kudo (1998) *Mol. Cell. Bioi.*, 18: 5492-5499). Therefore, we compare this pattern in CD43-positive and CD43-negative ovarian cell lines and tissue. Southern blot or PCR analysis is performed
5 in exactly the same way as described above with the exception that specific pairs of restriction endonucleases are employed. These pairs such as *MspI* and *HpaII* recognize the same nucleotide sequence but methylation of that sequence blocks digestion by one enzyme but not the other. Consequently, comparison by Southern blot or PCR analysis of the restriction patterns generated by digestion with each of such enzymes reflects the
10 methylation state of the CD43 gene.

Determination of Transcription Factor Structure in Ovarian Cancer. Irrespective of whether the CD43 gene present in ovarian cancer is amplified, mutant or wild-type, deletion and transfection studies are performed to delineate the region responsible for abnormal expression. EMSA, UV crosslinking and cloning are undertaken to characterize
15 the transcription factors with which this region interacts.

Transfection: In ovarian cancer cells where the CD43 gene is rearranged or contains point mutations, progressively smaller restriction fragments spanning these defects are isolated. Such fragments are then cloned in the appropriate orientation upstream of the promoterless luciferase gene present in the plasmid pATLuc (Shelley, *et al.*, (1993) *Proc. Natl. Acad. Sci. USA*, 90: 5364-5368). Three types of transfection are performed with the resulting constructs using methods we have previously described (Shelley, *et al.*, (1993) *Proc. Natl. Acad. Sci. USA*, 90: 5364-5368; Farokhzad *et al.*, (1996) *I. Immunol.*, 157: 5597-5605; Shelley *et al.*, (1991) *Proc. Natl. Acad. Sci. USA*, 88: 10525-10529; Shelley *et al.*, (1992) *Leukocyte adhesion* 337-351; Bottinger *et al.*, (1994)
25 *Mol. Cell. Biol.*, 14: 2604-2615). The first transfection is into ovarian cancer cell lines which express CD43 and the second is into primary cultures of ovarian tumors established on a fee-for-service basis by BioWhittaker, Inc. These first two transfection studies map to a resolution of approximately 100bp the region responsible for CD43 gene expression in ovarian cancer. The third set of transfections performed is into normal ovarian surface
30 epithelial cells immortalized by viral infection (Maines-Bandiera *et al.*, (1992) *Am. J. Obstet. Gynecol.*, 167: 729-725; Bruening *et al.*, (1999) *Cancer Res.*, 59: 4973-4983;

- 77 -

Gregoire *et al.*, (1998) *In Vitro Cell. Dev. Biol. Anim.*, 34: 636-639) and into primary cultures of normal ovarian cells purchased from BioWhittaker, Inc. These experiments establish if any identified mutation is solely responsible for driving abnormal expression of the CD43 gene in ovarian cells. As a negative control in all these studies mutant
5 constructs are transfected in parallel with p43B which contains the wild-type CD43 gene promoter.

If the CD43 gene is not mutated in ovarian cancer then the defects which cause abnormal CD43 expression may lie in *trans* to the CD43 gene. In order to induce transcription of the CD43 gene, ultimately these *trans-acting* defects must alter the nature
10 of transcription factor interaction with the CD43 gene. The region of the CD43 gene where these altered interactions occur is mapped by cloning progressively smaller fragments of the wild-type CD43 promoter into pATLuc. The resulting constructs is methylated *in vitro* to mimic any cancer specific methylation patterns detected by Southern blotting (Kudo *et al.*, (1995) *J. Biol. Chem.*, 270: 13298-13302) and transfected into
15 ovarian cell lines and primary cultures. In the event that primary ovarian cultures cannot be transfected either by electroporation or other methods such as those employing liposomes, calcium phosphate and DEAE dextran, virus infection is employed (Kolanus *et al.*, (1992) *EMBO J.*, 11: 4861-4868; Aftiring *et al.* (1994) *Hepatology*, 19: 1054-1056; Martiniello-Wilks *et al.*, (1998) *Hum. Gene Ther.*, 9: 1617-1626).

20 *EMSA*: The experiments outlined above define to a resolution of approximately 100bp the regions responsible for CD43 gene expression in ovarian cancer. A series of overlapping single and double stranded oligonucleotides covering these regions are synthesized and used in EMSA analyses to characterize the transcription factors with which they interact. Using methods we have previously described, nuclear extracts are
25 prepared from ovarian cancer cell lines and primary cultures as well as from immortalized and primary normal ovarian cultures (Shelley, *et al.*, (1993) *Proc. Natl. Acad. Sci. USA*, 90: 5364-5368; Farokhzad *et al.*, (1996) *I. Immunol.*, 157: 5597-5605; Shelley *et al.*, (1992) *Leukocyte adhesion* 337-351; Bottinger *et al.*, (1994) *Mol. Cell. Biol.*, 14: 2604-2615). These extracts are then incubated with the synthesized oligonucleotides
30 radiolabeled at their 5' ends. DNA-protein complexes are separated from unbound DNA by electrophoresis through native polyacrylamide gels and visualized by autoradiography.

- 78 -

Binding reactions are performed in the presence of non-radioactive competitor oligonucleotides that bind known transcription factors. In this way we determine the sequence specificity of the binding activities detected and the likelihood they represent factors already cloned. The identity of such factors is confirmed by antibody *Supershift* analyses performed as we have previously described (Shelley *et al.*, (1992) *Leukocyte adhesion* 337-351; Bottinger *et al.*, (1994) *Mol. Cell. Biol.*, 14: 2604-2615). DNA-protein complexes formed in nuclear extracts prepared from normal ovarian cells are compared to those formed in extracts of neoplastic ovarian cells to identify differences. Variations in the relative electrophoretic migration of complexes indicate qualitative differences while variations in their relative intensity, as assessed by densitometry, indicate quantitative differences. Once the CD43 gene is found to be mutated and/or abnormally methylated in ovarian cancers, the DNA-protein complexes which form on oligonucleotides containing these mutations and modifications are compared to those that form on oligonucleotides which do not.

UV Crosslinking: EMSA analyses can detect quantitative and gross qualitative differences in nuclear proteins, however, subtle qualitative differences are likely missed. We detect any such differences by determining the molecular weights of the nuclear proteins detected by EMSA (Farokhzad *et al.*, (1996) *J. Immunol.*, 157: 5597-5605; Pongubala *et al.*, (1992) *Mol. Cell. Biol.*, 12: 368-378). This we achieve using the technique of UV crosslinking as we have previously described (Farokhzad *et al.*, (1996) *J. Immunol.*, 157: 5597-5605). Single or double stranded oligonucleotides are synthesized with 5'-bromodeoxyuracil, radiolabeled and binding reactions performed as described for EMSA. After electrophoresis, protein-DNA complexes are irradiated *in situ* with UV light and visualized by autoradiography. Complexes excised from the gel are electroeluted, precipitated and subjected to SDS-PAGE in parallel with molecular weight markers. Autoradiography is performed again and the sizes of the DNA-binding proteins calculated by subtracting the weight of the oligonucleotide probes. The molecular weights of nuclear proteins expressed in normal and malignant ovarian cells is then compared for differences.

Cloning and Sequencing: The combination of transfection, EMSA, and UV crosslinking analyses described above characterize the cis-acting elements and *trans-acting* factors responsible for abnormal CD43 gene expression in ovarian cancer. These

- 79 -

analyses also determine which *trans*- acting factors exhibit abnormalities. Such abnormalities are identified by cloning and sequencing.

Factors already known to bind the CD43 gene, such as Sp1, hnRNP-K, Pura; and MeCP2, identified by EMSA will be cloned by reverse transcriptase (RT) PCR using RNA
5 isolated from ovarian cancer cell lines. RT-PCR is also sensitive enough to clone these factors from CD43-positive tumor cells isolated by LCM. The sequences of clones derived from malignant cells are compared to the published wild-type sequences for the identification of mutations.

There are a number of methods to clone unknown transcription factors (Singh, *et al.*, (1988) *Cell*, 52: 415-423; Wang *et al.*, (1993) *Nature (London)*, 364: 121-126; Hoey *et al.*, (1993) *Cell*, 72: 247-260). However, the most reliable involves classic protein
10 purification and limited sequencing followed by oligonucleotide screening (Hoey *et al.*, (1993) *Cell*, 72: 247-260). Putative transcription factors are purified from malignant and normal ovarian cell cultures using several chromatographic steps that include ion exchange
15 chromatography, fractionation on heparin agarose and ssDNA-sepharose. Finally, affinity chromatography is performed on a sequence-specific DNA resin consisting of 5' biotinylated binding-site multimers coupled to streptavidin agarose beads. Purification is followed throughout by EMSA and SDS-PAGE. If multiple bands are seen, each is electroeluted from the gel, renatured (Hager *et al.*, (1980) *Anal. Biochem.*, 109: 76-86)
20 and the gel retardation pattern, produced by each band alone or in combination, determined. Competitions using wild-type and mutant oligonucleotides are used to determine specificity in each case. The appropriate band is then cleaved with trypsin and the amino acid sequences of the resulting peptides determined as we have previously described using a LCQ Decca XP Plus electrospray mass spectroscope (ThermoFinnigan
25 Corp., San Jose, CA) set at MS/MS tandem mode (Da Silva *et al.*, (2002) *Blood*, 100: 3536-3544). Degenerate oligonucleotides is then synthesized based on peptide sequences and used to screen appropriate cDNA libraries. Full-length cDNA inserts are cloned into plasmid vectors which direct *in vitro* transcription and translation. The sizes of the translated products are then estimated by SDS-PAGE and compared to those of the
30 endogenous proteins as determined by UV crosslinking. The ability of the translated products to form complexes when mixed with wild-type or mutant binding-site

- 80 -

oligonucleotides is assessed using EMSA. The nucleotide sequences of clones derived from normal and malignant ovarian cells is compared to identify mutations. Next, factors containing any such mutations are transfected into cultures of normal ovarian cells to determine whether they are capable of activating the CD43 gene.

- 81 -

CLAIMS

1. A method for characterizing an ovarian cell, the method comprising:
determining the presence or absence of a CD43 molecule in an ovarian cell of a
5 subject to characterize the ovarian cell.
2. The method of claim 1, wherein the method is diagnostic and wherein the presence
of a CD43 molecule in the ovarian cell indicates that the subject has an ovarian tumor.
- 10 3. The method of claim 1, wherein the method is prognostic.
4. The method of claim 1, wherein the CD43 molecule is a CD43 nucleic acid.
5. The method of claim 1, wherein the CD43 molecule is a CD43 protein.
- 15 6. The method of claim 1, wherein determining the presence or absence of a CD43
molecule is performed *in vivo*.
7. The method of claim 1, wherein determining the presence or absence of a CD43
20 molecule is performed *in vitro*.
8. The method of any one of claims 1-7, wherein determining the presence or absence
of a CD43 molecule is done in the presence of a CD43 binding molecule.
- 25 9. The method of claim 8, wherein the CD43 binding molecule is a CD43 antibody.
10. The method of claim 9, wherein the CD43 antibody thereof is bound to a label.
11. The method of claim 10, wherein the label is selected from the group consisting of
30 a fluorescent label, an enzyme label, a radioactive label, a nuclear magnetic resonance
active label, a luminescent label, and a chromophore label.

- 82 -

12. The method of claim 8, wherein the CD43 binding molecule is an antigen-binding fragment of a CD43 antibody.

13. The method of claim 9, wherein the CD43 antibody is a CD43 monoclonal
5 antibody.

14. The method of claim 13, wherein the CD43 monoclonal antibody is selected from the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78,
10 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

15. A method of treating a subject having or at risk of having a tumor, comprising:
administering to a subject in need of such treatment one or more CD43 inhibitors in
an effective amount to treat the tumor.

15

16. The method of claim 15, wherein the CD43 inhibitor is a CD43 nucleic acid binding molecule.

17. The method of claim 16, wherein the CD43 nucleic acid binding molecule is an
20 hnRNP-K molecule.

18. The method of claim 17, wherein the hnRNP-K molecule is a hnRNP-K nucleic acid.

19. The method of claim 17, wherein the hnRNP-K molecule is a hnRNP-K protein.
25

20. The method of claim 15, wherein the CD43 inhibitor is transcription factor

21. The method of claim 20, wherein the transcription factor is a Pur α molecule.

30

22. The method of claim 21, wherein the Pur α molecule is a Pur α nucleic acid.

- 83 -

23. The method of claim 21, wherein the Pur α molecule is a Pur α protein.

24. The method of claim 15, wherein the CD43 inhibitor is a CD43 antisense molecule.

5 25. The method of claim 15, wherein the CD43 inhibitor is a CD43 antibody.

26. The method of claim 25, wherein the CD43 antibody is bound to a radioisotope.

27. The method of claim 26, wherein the radioisotope emits α radiations.

10

28. The method of claim 26, wherein the radioisotope emits β radiations.

29. The method of claim 26, wherein the radioisotope emits γ radiations.

15 30. The method of claim 26, wherein the radioisotope is selected from the group consisting of ^{225}Ac , ^{211}At , ^{212}Bi , ^{213}Bi , ^{186}Rh , ^{188}Rh , ^{177}Lu , ^{90}Y , ^{131}I or ^{67}Cu , ^{125}I , ^{123}I , and ^{77}Br .

31. The method of claim 25, wherein the CD43 antibody is bound to a therapeutic
20 moiety.

32. The method of claim 31, wherein the therapeutic moiety is a drug.

33. The method of claim 32, wherein the drug is a cytotoxic drug.

25

34. The method of claim 33, wherein the cytotoxic drug is selected from the group consisting of: calicheamicin, esperamicin, methotrexate, doxorubicin, melphalan, chlorambucil, ARA-C, vindesine, mitomycin C, cis-platinum, etoposide, bleomycin, and 5-fluorouracil.

30

- 84 -

35. The method of claim 31, wherein the therapeutic moiety is a toxin or a fragment thereof.

36. The method of claim 31, wherein the therapeutic moiety is an enzyme or a
5 fragment thereof.

37. The method of claim 25, wherein the CD43 antibody is a CD43 polyclonal antibody.

10 38. The method of claim 25, wherein the CD43 antibody is a CD43 monoclonal antibody.

39. The method of claim 38, wherein the CD43 monoclonal antibody is selected from the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14,
15 T2/53, 1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, .
148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

40. The method of claim 15, wherein the tumor is a solid tumor selected from the group consisting of: biliary tract cancer, brain cancer (including glioblastomas and
20 medulloblastomas), breast cancer, cervical cancer, choriocarcinoma, colon cancer, endometrial cancer, esophageal cancer, gastric cancer, intraepithelial neoplasms, including Bowen's disease and Paget's disease, liver cancer, lung cancer, lymphomas, including Hodgkin's disease and lymphocytic lymphomas, neuroblastomas, oral cancer, including squamous cell carcinoma, ovarian cancer, including those arising from epithelial cells,
25 stromal cells, germ cells and mesenchymal cells, pancreatic cancer, prostate cancer, rectal cancer, renal cancer including adenocarcinoma and Wilms tumor, sarcomas, including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma and osteosarcoma, skin cancer, including melanoma, Kaposi's sarcoma, basocellular cancer and squamous cell cancer, testicular cancer, including germinal tumors (seminomas, and non-seminomas such
30 as teratomas and choriocarcinomas), stromal tumors and germ cell tumors, and thyroid cancer, including thyroid adenocarcinoma and medullary carcinoma.

- 85 -

41. The method of claim 15, wherein the tumor is a non-solid tumor selected from the group of hematological neoplasms including: acute or chronic lymphocytic and myelogenous leukemia, multiple myeloma, AIDS associated leukemias and adult T-cell lymphoma/leukemia.

5

42. The method of claim 15, wherein the tumor is ovarian cancer.

43. The method of any one of claims 15-42, further comprising administering one or more anti-tumor therapy.

10

44. The method of claim 43, wherein the anti-tumor therapy comprises surgery, radiation therapy, or a chemotherapeutic agent.

15

45. The method of claim 43, wherein the anti-tumor therapy comprises an anti-ovarian tumor therapy.

46. The method of claim 45, wherein the anti-ovarian tumor therapy is an anti-ovarian tumor chemotherapeutic agent.

20

47. The method of claim 46, wherein the anti-ovarian tumor chemotherapeutic agent comprises cisplatin (Platinol) or platinum-containing drug combinations.

25

48. The method of claim 46, wherein the anti-ovarian tumor chemotherapeutic agent comprises one or more agents selected from the group consisting of: vinblastin, bleomycin, etoposide, hexamethylmelanine, gemcitabine, topotecan, ifosfamide, alkylating agents, progestational agents, and antiestrogens.

30

49. The method of claim 48, wherein the alkylating agent is selected from the group consisting of: Mechlorethamine (nitrogen mustard; HN2; Mustargen), Chlorambucil (Leukeran), Cyclophosphamide (Cytoxan), Melphalan (Alkeran), Thiotepa (triethylenethiophosphoramide), Busulfan (Myleran), Carmustine (BCNU), Lomustine

- 86 -

(CCNU), Semustine (methyl-CCNU), Altretamine (hexamethyl-melamine), Procarbazine (Matulane), Dacarbazine, and Carboplatin (Paraplatin).

50. The method of claim 48, wherein the progestational agent is selected from the group consisting of: progesterone, progesterone derivative, 17-ethinyl testosterone derivative, and 19-nortestosterone derivative.

51. The method of claim 48, wherein the progesterone derivative is selected from the group consisting of: hydroxyprogesterone caproate, medroxyprogesterone acetate, and megestrol acetate.

52. The method of claim 50, wherein the 17-ethinyl testosterone derivative is Dimethisterone.

53. The method of claim 50, wherein the 19-nortestosterone derivative is selected from the group consisting of: Desogestrel, Norethynodrel, Lynestrenol, Norethindrone, Norethindrone acetate, Ethynodiol diacetate, and L-Norgestrel.

54. The method of claim 48, wherein the antiestrogen is selected from the group consisting of: Tamoxifen, anastrozole, and a combination of aminoglutethimide and hydrocortisone.

55. The method of claim 15, wherein the ovarian tumor is selected from the group consisting of: serous cystoma, mucinous cystoma, endometrioid tumor, mesonephric tumor, transitional cell (Brenner) tumor, dermoid tumor, teratoma, fibroma, thecoma, luteoma, granulosa cell tumor, struma ovarii, dysgerminoma, malignant mixed mesodermal tumor, and carcinoma.

56. The method of claim 55, wherein the serous cystoma is a serous benign cystadenoma, a serous cystadenoma, or a serous cystadenocarcinoma.

- 87 -

57. The method of claim 55, wherein the mucinous cystoma is a mucinous benign cystadenoma, a mucinous cystadenoma, or a mucinous cystadenocarcinoma.

58. The method of claim 55, wherein the endometrioid tumor is an endometrioid
5 benign cyst, or an endometrioid adenocarcinoma.

59. The method of claim 55, wherein the mesonephric tumor is a benign mesonephric tumor, or a mesonephric cystadenocarcinoma.

10 60. The method of claim 55, wherein the carcinoma is a clear cell carcinoma.

61. The method of claim 15, wherein the ovarian tumor has metastasized beyond the ovary.

15 62. A method for assessing the regression or progression of an ovarian tumor in a subject with comprising the steps of:
measuring in a first ovarian cell obtained from the subject, the presence of a CD43 molecule,

measuring in a second ovarian cell obtained from the subject, the presence of a
20 CD43 molecule,

comparing the presence of the CD43 molecule in the first ovarian cell and the second ovarian cell,

wherein a decrease in the presence of the CD43 molecule in the second ovarian cell compared to the first ovarian cell indicates regression of the ovarian tumor, and

25 wherein an increase in the presence of the CD43 molecule in the second ovarian cell compared to the first ovarian cell indicates progression of the ovarian tumor.

63. The method of claim 62, wherein the CD43 molecule is a CD43 nucleic acid.

30 64. The method of claim 62, wherein the CD43 molecule is a CD43 protein.

- 88 -

65. The method of claim 62, wherein measuring the presence of a CD43 molecule is performed *in vivo*.

5 66. The method of claim 62, wherein measuring the presence of a CD43 molecule is performed *in vitro*.

67. The method of any one of claims 62-66, wherein measuring the presence of a CD43 molecule is done in the presence of a CD43 binding molecule.

10 68. The method of claim 67, wherein the CD43 binding molecule is a CD43 antibody.

69. The method of claim 68, wherein the CD43 antibody is a CD43 polyclonal antibody.

15 70. The method of claim 68, wherein the CD43 antibody is a CD43 monoclonal antibody.

71. The method of claim 70, wherein the CD43 monoclonal antibody is selected from the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14,
20 T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

72. A kit for diagnosing an ovarian tumor, comprising:
one or more CD43 binding molecules,
25 one or more control agents, and
instructions for the use of the CD43 binding molecules, and the control agents in the diagnosis of an ovarian tumor.

73. The kit of claim 72, wherein the CD43 binding molecule binds a CD43 nucleic acid.
30

- 89 -

74. The kit of claim 72, wherein the CD43 binding molecule binds a CD43 protein.

75. The kit of claim 72, wherein diagnosing an ovarian tumor is performed *in vivo*.

5 76. The kit of claim 72, wherein diagnosing an ovarian tumor is performed *in vitro*.

77. The kit of claim 72, wherein the CD43 binding molecule is a CD43 antibody.

78. The kit of claim 77, wherein the CD43 antibody is a CD43 polyclonal antibody.

10

79. The kit of claim 77, wherein the CD43 antibody is a CD43 monoclonal antibody.

80. The kit of claim 79, wherein the CD43 monoclonal antibody is selected from the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14,

15 T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

81. The kit of claim 72, wherein the one or more control agents are bound to a substrate.

20

82. A method of reducing the number of white blood cells in a subject comprising administering to a subject in need of a reduction in the number of white blood cells, a CD43 inhibitor in an effective amount to reduce the number of white blood cells in said subject.

25

83. The method of claim 82, wherein the CD43 inhibitor is a hnRNP-K molecule.

84. The method of claim 83, wherein the hnRNP-K molecule is a hnRNP-K nucleic acid.

30

85. The method of claim 83, wherein the hnRNP-K molecule is a hnRNP-K protein.

- 90 -

86. The method of claim 82, wherein the CD43 inhibitor is transcription factor.

87. The method of claim 86, wherein the transcription factor is a Pura molecule.

5

88. The method of claim 87, wherein the Pura molecule is a Pura nucleic acid.

89. The method of claim 87, wherein the Pura molecule is a Pura protein.

10 90. The method of claim 82, wherein the CD43 inhibitor is a CD43 antisense molecule.

91. The method of claim 82, wherein the CD43 inhibitor is a CD43 antibody.

15 92. The method of claim 91, wherein the CD43 antibody is a CD43 polyclonal antibody.

93. The method of claim 91, wherein the antibody is a CD43 monoclonal antibody.

20 94. The method of claim 93, wherein the CD43 monoclonal antibody is selected from the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

25 95. The method of claim 82, wherein the subject has a white blood cell tumor selected from the group consisting of: acute leukemias, chronic leukemias, and lymphomas.

96. The method of claim 82, wherein the subject is about to undergo, is undergoing, or has undergone a bone marrow, organ, cellular, or material transplant.

30 97. The method of any of claims 82-96, further comprising administering a second therapy to reduce the number of white blood cells.

- 91 -

98. A method of inhibiting a CD promoter comprising contacting the CD promoter with a CD43 inhibitor in an amount effective to inhibit the CD promoter.
- 5 99. The method of claim 98, wherein the CD43 inhibitor is a hnRNP-K molecule.
100. The method of claim 99, wherein the hnRNP-K molecule is a hnRNP-K nucleic acid.
- 10 101. The method of claim 99, wherein the hnRNP-K molecule is a hnRNP-K protein.
102. The method of claim 98, wherein the CD43 inhibitor is transcription factor.
103. The method of claim 102, wherein the transcription factor is a Pur α molecule.
- 15 104. The method of claim 103, wherein the Pur α molecule is a Pur α nucleic acid.
105. The method of claim 103, wherein the Pur α molecule is a Pur α protein.
- 20 106. The method of claim 98, wherein the CD43 inhibitor is a CD43 antisense molecule.
107. The method of claim 98, wherein the CD43 inhibitor is a CD43 antibody.
108. The method of claim 107, wherein the CD43 antibody is a CD43 polyclonal
- 25 antibody.
109. The method of claim 107, wherein the antibody is a CD43 monoclonal antibody.
110. The method of claim 109, wherein the CD43 monoclonal antibody is selected from
- 30 the group consisting of: BS1, MEM-59, 84-3C1, Bra7G, DF-T1, 1G10, MT1, L10, L14,

- 92 -

T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161.

111. The method of claim 98, further comprising administering a Pur α molecule.

5

112. The method of claim 98, wherein the CD promoter is inhibited *in vivo*.

113. The method of claim 98, wherein the CD promoter is inhibited *in vitro*.

10 114. The method of claim 98, wherein the CD promoter is selected from the group consisting of: CD43, CD11a, CD11b, CD11c, and CD11d.





Figure 1

6/10/2004 2:44 PM


JP14 Rec'd 23 MAR 2005

THIS PAGE BLANK (USPTO)

**Figure 2**

JC14 REG'D PGM 23 MAR 2005

THIS PAGE BLANK (USPTO)

**Figure 3**

JC14 Rec'd PCT/P 23 MAR 2005

THIS PAGE BLANK (USPTO)





Figure 4

2014 DEC 01 09:23 MAR 2005

THIS PAGE BLANK (USPTO)

**Figure 5**

JG14 Rec'd PGT/F 23 MAR 2005

THIS PAGE BLANK (USPTO)




Figure 6

JC14 Rec'd PCT/P 23 MAR 2005

THIS PAGE BLANK (USPTO)





Figure 7

2004/02/10 10:00:00

32 MAY 2005

THIS PAGE BLANK (USPT.O)

**Figure 8**

23 MAR 2005

THIS PAGE BLANK (USPTO)




Figure 9

J014 Rec'd PCI/P 23 MAR 2005

THIS PAGE BLANK (USPTO)




Figure 10

01/02/04 10:00 AM 10/528948

JC14 Rec'd PCT/

MAR 2005

THIS PAGE BLANK (USPTO)

SEQUENCE LISTING

<110> THE GENERAL HOSPITAL CORPORATION
 SHELLEY, CARL SIMON
 FAROKHZAD, OMID C.

<120> METHODS FOR DIAGNOSING AND TREATING TUMORS AND SUPPRESSING CD
 PROMOTERS

<130> M00765.70064

<140> not yet assigned

<141> 2003-09-23

<150> US 60/412,964

<151> 2002-09-23

<160> 28

<170> PatentIn version 3.2

<210> 1

<211> 1879

<212> DNA

<213> Homo sapiens sialophorin

<400> 1

gcctcgggag gtggtggagt gacctggccc cagtgtgcg tccttatcag ccgagcoggt	60
cccagctctt gctcctgcct gtttgcttg aaatggccac gcttctcctt ctccttgggg	120
tgctggtggt aagcccagac gctctgggga gcacaacagc agtgcagaca cccacctccg	180
gagagccttt ggtctctact agcagacccc tgagctcaaa gatgtacacc acttcaataa	240
caagtgaccc taaggccgac agcactgggg accagacctc agccctacct cctcgaactt	300
ccatcaatga gggatcccct ctttggaactt ccattggtgc cagcactggt tcccccttac	360
ctgagccaac aacctaccag gaagtttcca tcaagatgtc atcagtgcgc caggaaaccc	420
ctcatgcaac cagtcatcct gctgttccca taacagcaaa ctctctagga tcccacaccg	480
tgacagggtg aaccataaca acgaactctc cagaaacctc cagtaggacc agtggagccc	540
ctgttaccac ggcagctagc tctctggaga cctccagagg cacctctgga cccctctta	600
ccatggcaac tgtctctctg gagacttcca aaggcacctc tggaccccct gttaccatgg	660
caactgactc tctggagacc tccactggga ccaactggacc cctgtacc atgacaactg	720
gctctctgga gccctccagc ggggccagtg gacccaggt ctctagcgta aaactatcta	780
caatgatgtc tccaacgacc tccaccaacg caagcactgt gcccttccgg aaccagatg	840
agaactcacg aggcattgct ccagtggctg tgcttggtgc cctgctggcg gtcattagtc	900
togtggctct gctcctgctg tggcgccggc ggcagaagcg gcggactggg gccctcgtgc	960
tgagcagagg tggcaagcgt aacgggggtg tggacgcctg ggctgggcca gccaggtcc	1020

```

ctgaggaggg ggccgtgaca gtgaccgtgg gaggggtccgg gggcgacaag ggctctgggt 1080
tccccgatgg ggaggggtct agccgtcggc ccacgctcac cactttcttt ggcagacgga 1140
agtctcgcca gggctccctg gcgatggagg agctgaagtc tgggtcaggc cccagcctca 1200
aaggggagga ggagccactg gtggccagtg aggatggggc tgtggacgcc ccagctcctg 1260
atgagcccga agggggagac ggggctgcc ctttaagtgtc ggtgaatagt gaggctggag 1320
gccggaatct cagccagcct ccagcacctt ccctctcacc atcccactgc cccctcgctc 1380
ccatgtttcc acccggcacc ctgatcctca cccgaatctc cttttttttt ttcttttgag 1440
acagagtttc gctttgtcgc ccaggctgga gtgcaatgca cgatctcagt tcaactgcaac 1500
ctctgcctcc taagttcagg cgattctcct gcctcagctt cccgagtaac tgagattaca 1560
ggcacccacc accatgccca gctgcttttt tgtatttttg gtagagatgg ggtttcacca 1620
tgtttggctag gctggtctca aactcctgac ctcaggatgat ctacctgcct cagcctccca 1680
aagtgctgag attacagaca tgagcctccg cgccttgccct cctcaccac ctcttcaactc 1740
tgaatcctca tgaggcttct cagccctgga tttcctgctg ccacctctac ccagcaccca 1800
caactagcgc ctgggcaggg cagggtggc acctctcaac gtctgtggac tgaatgaata 1860
aaccctctc atccacccc 1879

```

<210> 2
 <211> 400
 <212> PRT
 <213> Homo sapiens sialophorin

<400> 2
 Met Ala Thr Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
 1 5 10 15
 Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30
 Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser

130. 135 140
 Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
 145 150 155 160
 Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
 165 170 175
 Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
 180 185 190
 Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
 195 200 205
 Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
 210 215 220
 Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
 225 230 235 240
 Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
 245 250 255
 Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
 260 265 270
 Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
 275 280 285
 Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
 290 295 300
 Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
 305 310 315 320
 Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
 325 330 335
 Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg
 340 345 350
 Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
 355 360 365
 Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
 370 375 380
 Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
 385 390 395 400

<210> 3
 <211> 1893
 <212> DNA
 <213> Homo sapiens

<400> 3
 cttccctgcc tccctcaggt cccagctctt gctcctgcct gtttgcttgg aaatggccac 60
 gcttctcctt ctccttgggg tgctggtggt aagcccagac gctctgggga gcacaacagc 120
 agtgcagaca cccacctccg gagagccttt ggtctctact agcgagcccc tgagctcaaa 180

gatgtacacc acttcaataa caagtgaccc taaggccgac agcactgggg accagacctc 240
agccctacct ccctcaactt ccatcaatga gggatcccct ctttggactt ccattgggtgc 300
cagcactgggt tccccctttac ctgagccaac aacctaccag gaagtttcca tcaagatgtc 360
atcagtgtccc caggaaaccc ctcatgcaac cagtcatcct gctgttccca taacagcaaa 420
ctctctagga tcccacaccg tgacaggtgg aaccataaca acgaactctc cagaaacctc 480
cagtaggacc agtggagccc ctgttaccac ggcagctagc tctctggaga cctccagagg 540
cacctctgga ccccccttta ccatggcaac tgtctctctg gagacttcca aaggcacctc 600
tggacccctt gttaccatgg caactgactc tctggagacc tccactggga cactggacc 660
ccctgttacc atgacaactg gctctctgga gccctccagc ggggccagtg gacccaggt 720
ctctagcgta aaactatcta caatgatgtc tccaacgacc tccaccaacg caagcactgt 780
gcccttccgg aaccagatg agaactcacg aggcagtctg ccagtggctg tgcttgtggc 840
cctgctggcg gtcatagtcc tcgtggctct gctcctgctg tggcgccggc ggcagaagcg 900
gcgactggg gccctcgtgc tgagcagagg cggcaagcgt aacggggtgg tggacgcctg 960
ggctgggcca gccaggtcc ctgaggagg ggcctgaca gtgacgtgg gaggtccgg 1020
gggcgacaag ggtctgggt tccccgatgg ggaggggtct agccgtcggc ccacgctcac 1080
cactttcttt ggcagacgga agtctcgcca gggctccctg gcgatggagg agctgaagtc 1140
tgggtcaggc ccagcctca aaggggagga ggagccactg gtggccagtg aggatggggc 1200
tgtggacgcc ccagctcctg atgagccga agggggagac ggggctgccc cttaagtgtc 1260
ggtgaatagt gaggtggag gccggaatct cagccagcct ccagcacctt ccctctcacc 1320
atcccactgc cccctcgtc ccatgtttcc acccggcacc ctgatcctca cccgaatctc 1380
cttttttttt ttcttttgag acagagtttc gctttgtcgc ccaggctgga gtgcaatgca 1440
cgatctcagt tcaactgcaac ctctgcctcc taagttcagg cgattctcct gcctcagctt 1500
cccagtaac tgagattaca ggcaccacc accatgccca gctgcttttt tgtatttttg 1560
gtagagatgg ggtttacca tgttggctag gctggctctc aactcctgac ctgaggtgat 1620
ctacctgcct cagcctccca aagtgtgag attacagaca tgagcctccg cgccttgctt 1680
cctcaccac ctcttcactc tgaatcctca tgaggcttct cagccctgga tttcctgctg 1740
ccatcctcac ccagcaccca caactagcgc ctgggcaggg cagggtggc acctctcaac 1800
gtctgtggac tgaatgaata aaccctctc atccaccctt aaaaaaaaaa aaaaaaaaaa 1860
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 1893

<210> 4
<211> 400
<212> PRT

<213> Homo sapiens

<400> 4

Met Ala Thr Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
1 5 10 15

Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
20 25 30

Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
35 40 45

Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
50 55 60

Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
65 70 75 80

Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
85 90 95

Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
100 105 110

Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
115 120 125

Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser
130 135 140

Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
145 150 155 160

Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
165 170 175

Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
180 185 190

Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
195 200 205

Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
210 215 220

Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
225 230 235 240

Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
245 250 255

Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
260 265 270

Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
275 280 285

Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
290 295 300

Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
305 310 315 320

Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
325 330 335

Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg
340 345 350

Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
355 360 365

Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
370 375 380

Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
385 390 395 400

<210> 5
<211> 1924
<212> DNA
<213> Human leukosialin

<400> 5
cctctgagcc cagccctccc tagcatcacc attccatcc cattcctcag ccaagagcca 60
ggaatcctga ttccagatcc cacgcttccc tgcctccctc aggtcccagc tcttgctcct 120
gcctgtttgc ctggaaatgg ccacgcttct ccttctcctt ggggtgctgg tggttaagccc 180
agacgctctg gggagcacia cagcagtgc gacacccacc tccggagagc ctttggtctc 240
tactagcgag cccctgagct caaagatgta caccacttca ataacaagt accctaaggc 300
cgacagcact ggggaccaga cctcagccct acctccctca attccatca atgagggatc 360
ccctctttgg atttcattg gtgccagcac tggttcccct ttacctgagc caacaacct 420
ccaggaagtt tccatcaaga tgtcatcagt gcccaggaa acccctcatg caaccagtca 480
tcctgctgtt ccataacag caaactctct aggatccac accgtgacag gtggaacct 540
aacaacgaac tctccagaaa cctccagtag gaccagtga gccctgtta ccacggcagc 600
tagctctctg gagacctcca gaggcacctc tggacccctt cttacctgag caactgtctc 660
tctggagact tccaaaggca cctctggacc ccdgtttacc atggcaactg actctctgga 720
gacctccact gggaccactg gacccctgt taccatgaca actggctctc tggagccctc 780
cagcggggcc agtggacccc aggtctctag cgtaaaacta tctacaatga tgtctccaac 840
gacctccacc aacgcaagca ctgtgccctt ccggaacca gatgagaact cagagggcat 900
gctgccagtg gctgtgcttg tggccctgct ggcggtcata gtccctcgtg ctctgctcct 960
gctgtggcgc cggcggcaga agcggcggac tggggccctc gtgctgagca gagcgggcaa 1020
gcgtaacggg gtgggtggacg cctgggctgg gccagcccag gtccctgagg agggggccgt 1080

gacagtgacc gtgggaggggt ccgggggcca caagggctct gggttccccg atggggaggg 1140
 gtctagccgt cggcccacgc tcaccacttt ctttggcaga cggaagtctc gccagggctc 1200
 cctggcgatg gaggagctga agtctgggtc agggcccagc ctcaaagggg aggaggagcc 1260
 actggtgggc agtgaggatg gggctgtgga cgcccagct cctgatgagc ccgaaggggg 1320
 agacggggct gccccttaag tgtcggtgaa tagtgaggct ggaggccgca atctcagcca 1380
 gcctccagca ccttccctct caccatccca ctgccccctc gctcccatgt ttccaccggg 1440
 caccctgac ctcaccgaa tctcctttt tttttcttt tgagacagag ttctgctttg 1500
 tcgcccaggc tggagtcaa tgcacgatct cagttcactg caacctctgc ctctaagtt 1560
 caggcgattc tcctgcctca gcttcccag taactgagat tacaggcacc caccaccatg 1620
 cccagctgct tttttgtatt tttggtagag atggggtttc accatgttgg ctaggctggt 1680
 ctcaaactcc tgacctcagg tgatctacct gcctcagcct cccaaagtgc tgagattaca 1740
 gacatgagcc tcgcgcctt gcctcctcac ccacctcttc actctgaatc ctcatgaggc 1800
 ttctcagccc tggatttctt gctgccatcc tcaccagca cccacaacta gcgcctgggc 1860
 agggcagggc tggcacctct caacgtctgt ggactgaatg aataaaccct cctcatccac 1920
 ccct 1924

<210> 6
 <211> 400
 <212> PRT
 <213> Homo sapiens leukosialin

<400> 6
 Met Ala Thr Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
 1 5 10 15
 Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30
 Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser

130	135	140
Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr		
145	150	155
Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu		
165	170	175
Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp		
180	185	190
Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr		
195	200	205
Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser		
210	215	220
Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala		
225	230	235
Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu		
245	250	255
Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala		
260	265	270
Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu		
275	280	285
Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala		
290	295	300
Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly		
305	310	315
Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser		
325	330	335
Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg		
340	345	350
Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser		
355	360	365
Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val		
370	375	380
Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro		
385	390	395
		400

<210> 7

<211> 2288

<212> DNA

<213> Homo spaiens leukosialin

<400> 7

ggagcctcgg gaggtggtgg agtgacctgg cccagtgct gcgtccttat cagccgagcc 60

ggtcccagct cttgctcctg cctgtttgcc tggaaatggc cagccttctc cttctccttg 120

gggtgctggt ggtaagccca gacgctctgg ggagcacaac agcagtgag acacccacct 180

ccggagagcc tttggtctct actagcgagc ccctgagctc aaagatgtac accacttcaa 240
taacaagtga ccctaaggcc gacagcactg gggaccagac ctgagcccta cctccctcaa 300
cttccatcaa tgagggatcc cctctttgga cttccattgg tgccagcact ggttccctt 360
tacctgagcc aacaacctac caggaagttt ccatcaagat gtcacagtg cccagggaaa 420
cccctcatgc aaccagtcac cctgctgttc ccataacagc aaactctcta ggatcccaca 480
ccgtgacagg tggaaccata acaacgaact ctccagaaac ctccagtagg accagtggag 540
cccctgttac cacggcagct agctctctgg agacctccag aggcacctct ggacccccctc 600
ttaccatggc aactgtctct ctggagactt ccaaaggcac ctctggaccc cctgtttacca 660
tggaactga ctctctggag acctccactg ggaccactgg accccctgtt accatgacaa 720
ctggctctct ggagccctcc agcggggcca gtggaccca ggtctctagc gtaaaactat 780
ctacaatgat gtctccaacg acctccacca acgcaagcac tgtgcccttc cggaaccocag 840
atgagaactc acgaggcatg ctgccagtgg ctgtgcttgt ggccctgctg gcggtcatag 900
tcctcgtggc tctgctcctg ctgtggcgcc ggccggcagaa gcggcggaact ggggccctcg 960
tgctgagcag aggcggcaag cgtaacgggg tggtagacgc ctgggctggg ccagcccagg 1020
tccctgagga gggggccgtg acagtgaccg tgggagggtc cgggggcgac aagggtctctg 1080
ggttccccga tggggagggg tctagccgtc ggccacgct caccacttct tttggcagac 1140
ggaagtctcg ccagggctcc ctggcgatgg aggagctgaa gtctgggtca ggccccagcc 1200
tcaaagggga ggaggagcca ctggtggcca gtgaggatgg ggctgtggac gccccagctc 1260
ctgatgagcc cgaaggggga gacggggctg ccccttaagt gtcggtgaat agtgaggctg 1320
gaggccgaa tctcagccag cctccagcac ctccctctc accatccac tgccccctcg 1380
ctcccatgtt tccaccggc acctgatcc tcaccogaat ctctttttt tttttcttt 1440
gagacagagt ttgccttgt cgccaggct ggagtgaat gcacgatctc agttcactgc 1500
aacctctgcc tcctaagttc aggcgattct cctgcctcag cttcccgagt aactgagatt 1560
acaggcacc accacatgc ccagctgctt ttttgtattt ttggtagaga tggggtttca 1620
ccatgttggc taggctggtc tcaaactcct gacctcaggat gatctacctg cctcagcctc 1680
ccaaagtgt gagattacag acatgagcct ccgcgccttg cctcctcacc cacctcttca 1740
ctctgaatcc tcatgaggct tctcagccct ggatttctct ctgccatct caccagcac 1800
ccacaactag cgcttgggca gggcagggtt ggcacctctc aacgtctgtg gactgaatga 1860
ataaacctc ctcttacaaa tgccaaaatt cattcagctt tgatgataaa cactgaggcc 1920
caatggcctt tatcatctag ggagtatgaa gaatgagcaa gaggctaact cagcgtgagt 1980
taccctggga aaggaaagaa gaaatggctc atataggaca cacatagata ccttcaaggg 2040

tgtttgtaga gttgtttcctt aagtagttgg ttctcttcac agaaagttct taaactcaga 2100
 atatacccat ccatgcaccc caccagcaat acaaagaccc caaacaacaaa attactatat 2160
 tcttacccta cgcattggctt cctcctcttc ttgacgcttt tcataatgtg caaagtcac 2220
 aaagattgag gtggtatgct tgaaagtagc aattatttta agcacttgct tagctttttc 2280
 aagggacg 2288

<210> 8
 <211> 400
 <212> PRT
 <213> Homo sapiens leukosialin

<400> 8
 Met Ala Thr Leu Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
 1 5 10 15
 Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30
 Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser
 130 135 140
 Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
 145 150 155 160
 Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
 165 170 175
 Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
 180 185 190
 Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
 195 200 205
 Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
 210 215 220
 Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
 225 230 235 240

Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
245 250 255

Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
260 265 270

Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
275 280 285

Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
290 295 300

Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
305 310 315 320

Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
325 330 335

Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg
340 345 350

Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
355 360 365

Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
370 375 380

Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
385 390 395 400

<210> 9

<211> 5050

<212> DNA

<213> Homo sapiens leukosialin (CD43)

<400> 9

```

ccccctgca gaatgggcac cccgttacct ttctgagcca ctgtgcgcag aaaagagagc      60
atgttggcca ggctggtctc gaactcctga cctcaagtga tcagcctgcc ttacctccca      120
aagtccctggg attacaggcg tgaaccacca cgctcagcct ctgaatactt tgtactcaag      180
ccatttttca gtgctgtgtt tgcagtgagc acaccgagg gatgaagaca cgtctccctg      240
tggaacctg ggcttaccag ggcccctaga ggaggggaat ctctcaagct cagagctcta      300
tggtgcggt gcaggccac tgtgtgcatg gtgtcagtct gggcccttcc atgttgcccc      360
cgtgggactt ggggtaaggg gaactgatgc aaacatcacg ctgctgttgc ttgggtgtgag      420
caattaattc ctgtggctct caccaggag tctcatgtct ttgggtcaga caaactcatc      480
agcttgtaga aatggcacag tcccacgggc ctgttagaat cttctattgt gcacatgttg      540
ctcttaaaat atacaaatca gttttgattt taaaaaatta tttatttttt tagtgatagg      600
agttttgcta cgttgcccag gctggtttca aactcttggg ctcaggaggt cctcccactt      660
tggcctggac tgccagcata atgtatcacc acaccggga ctgattttcg tttttcaaga      720

```

acaaaaacca aaaacataca caaaccgaga gtcaaagctt gctaattaga ggaaagtcag 780
gaaatgggaa ccattcaaag aagaaaatac cccacctcc tactctcacc tatccaaaga 840
caattaggtg aatcccttag tagatatctt tocagacggt tttccatata gattcccata 900
tctggccagg cgcggtggct cacacctgta atcctagcgc ttggggaggc tgaggcggat 960
ggaccacctg aggtcaggag ttcgagacca gcctgaccaa catggagaaa cctcgtctct 1020
acgaaaaata caaaattagc cgggcacagt ggtgcaagcc tgtaatcca gctactcagg 1080
aggccgaggc aggagaattg cttgaacctt ggaggcagac attgtgctga gccgagccaa 1140
gatcatgcca ttgcactaaa ctccgcctta aaaaaaaaaa aaaagattcc cacatcttta 1200
ctagtttgca gaaataagat cctagcatat gcagtgtgta ggaaccacct tggtttagcc 1260
acgtctctgt gactgggggc cactgtggtg acccccagct ccccgacag agtcaagagc 1320
tcaccagcct gcaaaggttt tcacggcccc cagccagact cgggggcttc ctcttgccct 1380
gctacttcct gggagctctg agggcaggaa atggcgccac tcagctcctg gcctaacagc 1440
ttggggacca caaatgcaaa ggaaaccacc ctcccctccc acctcctcct ctgcaccctt 1500
gagttctcag gctcacattc ccaccaccca cctctgagcc cagccctccc tagcatcacc 1560
acttccatcc cattcctcag ccaagagcca ggaatcctga ttccagatcc cacgcttccc 1620
tgctccctc aggtgagccc cagaccccca ggaccccgc tggccctga aggagcaggt 1680
gatggtgctg tcttcgcccc gcagctgtgg gacagggcgg gtggggcagg atggaggggt 1740
gggtggggtg ggtggagcca gggccactt cctttcccct tggggccctg tccttcccag 1800
tcttgcccc gctcgggag gtggtggagt gacctggccc cagtgtgcg tccttatcag 1860
ccgagccggt aagaggggtga gacttggtgg ggtaggggcc tcagtgggcc tgggaatgtg 1920
cctgtggctt gaaaagactc tgacaggtta tgatgggaag agattgggag ccattgggct 1980
gcacagggtc agggaaggcc aggaggggct ggtcactgct ggaatctaag ctgctgaggc 2040
tgaggggagc ctcaggatgg ggctgatgg ggagctgcca gcatctgttc ctctgtcatt 2100
tctgataaca gtaaaagcca gcatggaaaa aaccgttaaa ccgcagggtg ggcctggccg 2160
ttggcagggg agtgggcaga ggggaggccc ggccaggctc tccggcaact cccgcgtgtt 2220
ctgcttctcc ggctgcccac ctgcaggctc cagctcttgc tctgcctgt ttgcctggaa 2280
atggccacgc ttctccttct ccttgggggtg ctggtggtaa gccagacgc tctggggagc 2340
acaacagcag tgcagacacc cacctccgga gagcctttgg tctctactag cgagcccctg 2400
agctcaaaga tgtacaccac ttcaataaca agtgacctta aggccgacag cactggggac 2460
cagacctcag ccctacctcc ctcaacttcc atcaatgagg gatcccctct ttggacttcc 2520
attggtgcca gcactggttc ccctttacct gagccaacaa cctaccagga agtttccatc 2580

aagatgtcat cagtgtcccca ggaaaccct catgcaacca gtcacctgc tgttcccata 2640
acagcaaact ctctaggatc ccacaccgtg acaggtggaa ccataacaac gaactctcca 2700
gaaacctcca gtaggaccag tggagccct gttaccacgg cagctagctc tctggagacc 2760
tccagaggca cctctggacc ccctcttacc atggcaactg tctctctgga gacttccaaa 2820
ggcacctctg gacccctgt taccatggca actgactctc tggagacctc cactgggacc 2880
actggacccc ctgttaccat gacaactggc tctctggagc cctccagcgg ggccagtggga 2940
ccccaggtct ctagcgtaaa actatctaca atgatgtctc caacgacctc caccaacgca 3000
agcactgtgc ccttccggaa ccagatgag aactcacgag gcatgtgcc agtggctgtg 3060
cttgtggccc tgctggcggc catagtcctc gtggctctgc tctgtctgtg ggcggcgcg 3120
cagaagcggc ggactggggc cctcgtgctg agcagaggcg gcaagcgtaa cggggtgggtg 3180
gacgcctggg ctggggcagc ccaggctcct gaggaggggg ccgtgacagt gaccgtggga 3240
gggtccgggg ggcacaaggc ctctgggttc ccgatgggg aggggtctag ccgtcggccc 3300
acgctcacca ctttctttgg cagacggaag tctcgccagg gctccctggc gatggaggag 3360
ctgaagtctg ggtcaggccc cagcctcaaa ggggaggagg agccactggg ggccagtgag 3420
gatggggctg tggacgccc agctcctgat gagccgaag ggggagacgg ggctgccct 3480
taagtgtcgg tgaatagtga ggctggaggc cgcaatctca gccagcctcc agcaccttc 3540
ctctcaccat ccactgccc cctcgctccc atgtttccac ccggcaccct gatcctcacc 3600
cgaatctcct tttttttttt cttttgagac agagtctgc tttgtgccc aggctggagt 3660
gcaatgcaag atctcagtc actgcaacct ctgcctcta agttcaggcg attctcctgc 3720
ctcagcttcc cgagtaactg agattacagg caccaccac catgcccagc tgcttttttg 3780
tatttttggg agagatgggg ttccaccatg ttggctaggc tggctctcaa ctctgacct 3840
caggtgatct acctgcctca gcctcccaaa gtgctgagat tacagacatg agcctccgcg 3900
ccttgccctc tcaccacct cttaactctg aatcctcat aggtttctca gccctggatt 3960
tctgtctgcc atcctcacc agcaccaca actagcgctt ggcagggca gggctggcac 4020
ctctcaacgt ctgtggactg aatgaataaa cctcctcat ccaccctat ttatctccat 4080
caccatttcc ccctctttct tgttctgga aacggctgct gagtctccat cggccaaact 4140
tatctgcct gtgatttct tgacaattct ccttttcccc cagaaccac cctgggttga 4200
ccagagtctg ggaagaagga caagagaacc cggcaaactc cctcctagga ttaactttgt 4260
aaagcacct tgccctgtag ctgcaagggc tgtggaacct gggcagccc caaccacctt 4320
tagctctggg ccccccaggc cagcctggag catggctggg tggggccacc agcccatgct 4380
ctcaggcggg cctgtgatct ttcccagggc acatggactg taggctggcc ctggcccaca 4440

```

ccaccacact ctccccagcc atggacagag gcagccagag gcctcacggt ttctctccg 4500
agtttctggc tgggtgtagt tctcagaaac cccagtgcct gcgtgtgtcc actcgtgggt 4560
gtggtttgtg tgcaagagct gaggatttgg cgatgcttgg gaggggtagt tgtgggtaca 4620
gacgggtgtg ggggtgggaag tgggtgcagag actgaagagg gtcaacctgg gcatggggga 4680
cacaggggact gctgagaacg tgcgtgtcat ctttgctctg atgggggtgga catagcagaa 4740
aatctaactc tgtctgtagc cccatacaga atgccagggt gagcacagtg gctgggtgcct 4800
ttaatcccag cactttggaa agttgaggca ggaggatcgc ttgagcccag gaggttcgagt 4860
ctgaagtgag ctgtgattgc accactgcac ttcagcctgg gcaacagagt gagcccctgt 4920
ctcaaaaaag aaaagaaaaa gaaagccagg cttcatggaa agatcgtatg tgtgacccaa 4980
tatgagttct tcagctcagc catggtaatc ccttccttga agtctccatt tctgcagtac 5040
acatgcatgt 5050

```

<210> 10
 <211> 400
 <212> PRT
 <213> Homo sapiens leukosialin (CD43)

<400> 10
 Met Ala Thr Leu Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
 1 5 10 15
 Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30
 Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser
 130 135 140
 Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
 145 150 155 160
 Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
 165 170 175

Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
 180 185 190
 Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
 195 200 205
 Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
 210 215 220
 Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
 225 230 235 240
 Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
 245 250 255
 Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
 260 265 270
 Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
 275 280 285
 Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
 290 295 300
 Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
 305 310 315 320
 Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
 325 330 335
 Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg
 340 345 350
 Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
 355 360 365
 Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
 370 375 380
 Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
 385 390 395 400

<210> 11

<211> 1879

<212> DNA

<213> Homo sapiens sialophorin (CD43)

<400> 11

gcctcgggag gtggtggagt gacctggccc cagtgtctgcg tccttatcag ccgagccggt 60
 cccagctctt gctcctgcct gtttgectgg aaatggccac gcttctcctt ctccctgggg 120
 tgctgggtgt aagcccagac gctctgggga gcacaacagc agtgcagaca cccacctccg 180
 gagagccttt ggtctctact agcgagcccc tgagctcaaa gatgtacacc acttcaataa 240
 caagtgacct taaggccgac agcaactggg accagacctc agccctacct cctcaactt 300
 ccatcaatga gggatcccct ctttggaactt ccattgggtgc cagcaactgg tcccctttac 360

```

ctgagccaac aacctaccag gaagtttcca tcaagatgtc atcagtgtcc caggaaaccc 420
ctcatgcaac cagtcattct gctgttccca taacagcaaa ctctctagga tcccacaccg 480
tgacaggtgg aaccataaca acgaactctc cagaaacctc cagtaggacc agtggagccc 540
ctgttaccac ggcagctagc tctctggaga cctccagagg cacctctgga cccctcttta 600
ccatggcaac tgtctctctg gagacttcca aaggcaacctc tggacccctc gttaccatgg 660
caactgactc tctggagacc tccactggga ccaactggacc ccctgttacc atgacaactg 720
gctctctgga gccctccagc ggggccagtg gaccccaggt ctctagcgta aaactatcta 780
caatgatgtc tccaacgacc tccaccaacg caagcactgt gcccttcagg aaccagatg 840
agaactcacg aggcattgtg ccagtggctg tgcttgtggc cctgctggcg gtcattagtc 900
tcgtggctct gctcctgtg tggcgccggc ggcagaagcg gcggactggg gccctcgtgc 960
tgagcagagg tggcaagcgt aacgggggtg tggacgcctg ggctgggcca gccagggtcc 1020
ctgaggaggg ggccgtgaca gtgaccgtgg gagggctcgg gggcgacaag ggctctgggt 1080
tccccgatgg ggaggggtct agccgtcggc ccacgctcac cactttcttt ggagacgga 1140
agtctcgcca gggctccctg gcgatggagg agctgaagtc tgggtcaggc cccagcctca 1200
aaggggagga ggagccactg gtggccagtg aggatggggc tgtggacgcc ccagctcctg 1260
atgagccoga agggggagac ggggctgccc cttaagtgtc ggtgaatagt gaggtggag 1320
gccggaatct cagccagcct ccagcacctt ccctctcacc atccactgc ccctcgtctc 1380
ccatgtttcc acccggcacc ctgattctca cccgaatctc cttttttttt ttcttttgag 1440
acagagtttc gttttgtcgc ccaggctgga gtgcaatgca cgattctcagt tcaactgcaac 1500
ctctgcctcc taagttcagg cgattctcct gcctcagctt cccgagtaac tgagattaca 1560
ggcaccacc accatgccca gctgcttttt tgtatttttg gtagagatgg ggtttcacca 1620
tgttggttag gctggtctca aactcctgac ctcaggatgat ctacctgcct cagcctccca 1680
aagtgtgag attacagaca tgagcctccg cgcttgcct cctcaccac ctcttcaactc 1740
tgaatcctca tgaggcttct cagccctgga tttcctgctg ccatcctcac ccagcaccca 1800
caactagcgc ctgggcaggg cagggtggc acctctcaac gtctgtggac tgaatgaata 1860
aaccctctc atccacccc 1879

```

```

<210> 12
<211> 400
<212> PRT
<213> Homo sapiens sialophorin (CD43)

```

```

<400> 12
Met Ala Thr Leu Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
1           5           10           15

```

Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30
 Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser
 130 135 140
 Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
 145 150 155 160
 Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
 165 170 175
 Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
 180 185 190
 Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
 195 200 205
 Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
 210 215 220
 Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
 225 230 235 240
 Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
 245 250 255
 Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
 260 265 270
 Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
 275 280 285
 Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
 290 295 300
 Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
 305 310 315 320
 Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
 325 330 335
 Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg

340

345

350

Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
 355 360 365
 Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
 370 375 380
 Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
 385 390 395 400

<210> 13

<211> 6503

<212> DNA

<213> Homo sapiens sialophorin (CD43)

<400> 13

```

aagcttgcta attagaggaa agtcaggaaa tgggaaccat tcaaagaaga aaataccccc      60
acctcctact ctacactatc caaagacaat taggtgaatc ccttagtaga tatctttcca      120
gacgggttttc catatagatt cccatatctg gccaggcgcg gtggctcaca cctgtaatcc      180
tagcgcttgg ggaggctgag gcggatggac cacctgaggt caggagtctg agaccagcct      240
gaccaacatg gagaaacctc gtctctacga aaaatacaaa attagccggg cacagtgggtg      300
caagcctgta atcccagcta ctccaggaggc cgaggcagga gaattgcttg aacctaggag      360
gcagacattg tgctgagccg agccaagatc atgccattgc actaaactcc gccttaaaaa      420
aaaaaaaaaa agattccac atctttacta gtttgacagaa ataagatcct agcatatgca      480
gtgtgtagga accaccttgg tttagccacg tctctgtgac tgggggccac tgtggtgacc      540
cccagctccc cggacagagt caagagctca ccagcctgca aagggttttca cggccccag      600
ccagactcgg gggcttcctc ttgccctgct acttctctggg agctctgagg gcaggaaatg      660
gcgccactca gtccttgccc taacagcttg gggaccacaa atgcaaagga aaccaccctc      720
ccctcccacc tctcctctg cacccttgag ttctcaggct cacattccca ccaccacct      780
ctgagcccag cctcccttag catcaccact tccatcccat tctcagcca agagccagga      840
atcctgattc cagatccac gcttccctgc ctccctcagg tgagccccag acccccaggc      900
accccgctgg ccctgaagg agcaggatgat ggtgctgtct tcgcccagca gctgtgggag      960
caggcgggtg gggcaggatg gaggggtggg tggggtgggt ggagccaggg cccacttcct     1020
tcccccttgg ggcctgtcc ttcccagtct tgccccagcc tcgggaggtg gtggagtgcac     1080
ctggccccag tgetgcgtcc ttatcagccg agccggtgag aggggtgagac ttggtgggggt     1140
aggggcctca gtgggcctgg gaatgtgcct gtggcttgaa aagactctga caggttatga     1200
tggaagaga ttgggagcca ttgggctgca cagggtcagg gaaggccagg aggggctggt     1260
cactgctgga atctaagctg ctgaggctgg agggagcctc aggatggggc tgatggggga     1320
gctgccagca tctgttcctc tgtcatttct gataacagta aaagccagca tggaaaaaac     1380

```


cgtaaaccg caggttgggc ctggccgttg gcagggaagt gggcagaggg gagggccggc 1440
caggtcctcc ggcaactccc gcgtgttctg cttctccggc tgcccacctg caggtcccag 1500
ctcttgetcc tgctgtttg cctggaaatg gccacgcttc tcttctctct tggggtgctg 1560
gtggtaagcc cagacgctct ggggagcaca acagcagtgc agacaccac ctccggagag 1620
ccttttgtct ctactagcga gccctgagc tcaaagatgt acaccacttc aataacaagt 1680
gaccctaagg ccgacagcac tggggaccag acctcagccc tacctccctc aacttccatc 1740
aatgagggat cccctctttg gacttccatt ggtgccagca ctggttcccc ttacctgag 1800
ccaacaacct accaggaagt ttccatcaag atgtcatcag tgcccagga aacctctcat 1860
gcaaccagtc atctgtctgt tcccataaca gcaaactctc taggatcca caccgtgaca 1920
ggtggaacca taacaacgaa ctctccagaa acctccagta ggaccagtgg agcccctgtt 1980
accacggcag ctagctctct ggagacctcc agaggcacct ctggaccccc tcttaccatg 2040
gcaactgtct ctctggagac ttccaaaggc acctctggac cccctgttac catggcaact 2100
gactctctgg agacctcac tgggaccact ggaccccctg ttaccatgac aactggctct 2160
ctggagccct ccagcggggc cagtggacct caggtctcta gcgtaaaact atctacaatg 2220
atgtctccaa cgacctcac caacgcaagc actgtgccct tcgggaacct agatgagaac 2280
tcacgaggca tgctgccagt ggctgtgctt gtggccctgc tggcggtcat agtcctcgtg 2340
gctctgctcc tgctgtggcg ccggcggcag aagcggcgga ctggggccct cgtgctgagc 2400
agaggcggca agcgtaacgg ggtggtggac gcctgggctg ggccagccca ggtccctgag 2460
gagggggccg tgacagtgc cgtgggaggg tccggggcg acaagggtc tgggttcccc 2520
gatggggagg ggtctagccg tcggcccacg ctcaccactt tctttggcag acggaagtct 2580
cgccagggct ccttgccgat ggaggagctg aagtctgggt caggccccag cctcaaagg 2640
gaggaggagc cactggtggc cagtgaggat ggggctgtgg acgcccagc tctgatgag 2700
cccgaagggg gagacggggc tgccccttaa gtgtcggtga atagtaggc tggaggccgg 2760
aatctcagcc agcctccagc accttccctc tcaccatccc actgcccct cgtcccatg 2820
tttccaccgg gcacctgat cctcaccga atctcctttt ttttttctt ttgagacaga 2880
gtttcgcttt gtcgccagg ctggagtga atgcacgac tcagttcact gcaacctctg 2940
cctcctaagt tcaggcgatt ctctgcctc agcttccga gtaactgaga ttacaggcac 3000
ccaccacat gccagctgc tttttgtat ttttgtaga gatggggtt caccatgttg 3060
gctaggctgg tctcaaatc ctgacctcag gtgatctacc tgccctagcc tcccaaagt 3120
ctgagattac agacatgagc ctccgcgct tgccctccta cccacctct cactctgaat 3180
cctcatgagg cttctcagcc ctggatttcc tgctgccatc ctacccagc acccacaact 3240

agcgctggg cagggcaggg ctggcacctc tcaacgtctg tggactgaat gaataaaccc 3300
tcctcatcca cccctattta tctccatcac catttcccc tctttcttgt tcctggaaac 3360
ggctgctgag tctccatcgg ccaaacttat ctgccctgtg atttctttga caattctcct 3420
tttccccag aaccaccct gggttgacca gagtctggga agaaggacaa gagaaccgg 3480
caaactccct cctaggatta actttgtaaa gcacccttgc cctgtagctg caagggctgt 3540
ggaacctggg cagcccga ccaccttag ctctgggccccc ccaggccag cctggagcat 3600
ggctgggtgg ggccaccagc ccatgtctc aggcgggcct gtgatcttcc ccagggcaca 3660
tggactgtag gctggccctg gccacacca ccacactctc ccagccatg gacagaggca 3720
gccagaggcc tcacggttcc tcctccgagt ttctggctgg gtgtagttct cagaaacccc 3780
agtgcctgcg tgtgtccact cgtgggtgtg gtttgtgtgc aagagctgag gatttggcga 3840
tgcttgggag gggtagttgt gggtagacac ggtgtggggg tgggaagtgg tgcagagact 3900
gaagaggggc aacctgggca tgggggacac agggactgct gagaacgtgc gtgtcatctt 3960
tgctctgatg gggtagacat agcagaaaat ctaactctgt ctgtagcccc atacagaatg 4020
ccagggtag cacagtggct ggtgccttta atcccagcac tttggaaagt tgaggcagga 4080
ggatcgcttg agcccaggag ttcgagtctg aagttagctg tgattgcacc actgcacttc 4140
agcctgggca acagagttag cccctgtctc aaaaaagaaa agaaaaagaa agccaggctt 4200
catggaaaga tcgtatgtgt gacccaaata tgagttcttc agctcagcca tggtaatccc 4260
ttccttgaag tctccatttc tgcagtacac atgcatgtgc gctctctctc tctctctctc 4320
tctcacacac acacacacac acacacacac gcgcgcgcgc gcgcgcgcgc gcgctctcct 4380
gcgaacagag gcagggggag aggggttgc cctggtctcg gggactggtc tggctggcgc 4440
ttccccactg caggtttcca ggtttagttt gtctgtgtct cctcttccat ccaggggct 4500
gagccccttc catcctccaa gaggaaccag tgagagttag tgaaggagg gcctggagcc 4560
agggacttcc cctgtggggc ctgggtggag aggggagaa tcaatggtgc tgcctttgag 4620
accagcccag ctacagccca ggagcacaca tggggccagg cagttggtat ttcccagga 4680
caaagaggaa attttcaaag aggaagtgtg tgagttagag cttgcggtgg ctgagagcag 4740
acaggttgac ctgcaaaaaa agacagggga ggcatgtgag tgtgacagcc ctgctctgtg 4800
gcctgggcag gagatggggg aaagggtcag gtgggggatg ggctcgtgca gtgggagagg 4860
agacggaggg agggagcggg aaggggcttg cttagtgggt gggaagagct gagctcggat 4920
ggaaccagct tctaccagcc aggtgggca cccactgggc tgcactcgtt gcccttttct 4980
gattgctatt tggactcact gcagctgcag aatgacagag gccatgtcca aaatccctta 5040
gagacactgt tgtcttagag ttgttaaaat aagagccccc atatcagggt tagaaaatac 5100

tgtcaccgaa cgaacgtcgc tgtcctcagc tccacctccc tttcctttga cagatatggt 5160
 tgttttctaa gccaggactg gtttttagtca ggtcctgggc gaatcctgaa aaaaagaggt 5220
 agtacgggta aggaaggcac ccaacagggc tttcacaatc cagaaaatat caaaatataa 5280
 gtgttaaaag agaggcacag gccgggtgcg gtggctcacg cctgtaatct cagcactttg 5340
 ggaggccaag gtgggcagat catgaggta ggagtttgag accagcctgg ccaatatgat 5400
 gaaaccccgt ttctactaaa aatacaaaaag ttagccaggc atggtggtgt gtcctgttaa 5460
 tcccagctac ttaggaggct gaggccagag aattgcttga accctggagt cagagggttg 5520
 agtgagccgg gatcatgcc ctgtactcca ggctgggtga caaagtgaga ctgtctcaaa 5580
 aaataaaaat aaataaaaata aataaaaagag aggcacaaac agtgttatga atgcaccaag 5640
 gaaaatggtg cattcataac tctcagggtga agcctaccaa gccatgcgtg tgtgcacata 5700
 tgtgtgtacg tgtgcatgtg cgtgcgtgca tgtgcgtgcg tgcattgtgc tgtgtgtgta 5760
 tgtgtgcaca tgtgtgtgcg catgtgtgtg tgtgcgcgca tgtgtgtgtg catgcatgtt 5820
 ctcccatgca tgtgtactgt ggcaaggag actttgagga agagattcca gtggctgagc 5880
 agaagggtc gcattgccct ggcgaaaggt tggaaggctt cacctgagag tgtgtcgtgg 5940
 cctttgtcat atccactgct tgattccttt ctttaaaaat tatttttatt gttttctaca 6000
 tatgagaacc accacacctg gctaattttt gtattttttg tagagatggg gtttcacat 6060
 gttgtcccg ctggtctcaa actcccgggc acaagagatc cacctgcctc agcctcccaa 6120
 aatgctggga ctataggcat gagccactgc accagccac tgcttcattc ctggtggctg 6180
 ctgtgcctgg catgttgag atcctccatg aatatgcatt tgaatgaatg aatgaatgaa 6240
 tgaatgaatg gagatgacgc ctcagagatt ctttcttttg agatgaggtc tcattctgtc 6300
 acccagacta gagggcagtg gtgcaatcac agctcaccac agcctcaacc tcctgggctc 6360
 cccaagtagc tgcgatcaca ggtgtgcacc aacatgccca gctaattttt tttttaattt 6420
 ttaatttgta cagacagggt cttgctgtgt tgcccaggct ggtctcgaac tcctgggctc 6480
 aagtggctcct cccacctaag ctt 6503

<210> 14
 <211> 400
 <212> PRT
 <213> Homo sapiens sialophorin (CD43)

<400> 14
 Met Ala Thr Leu Leu Leu Leu Gly Val Leu Val Val Ser Pro Asp
 1 5 10 15
 Ala Leu Gly Ser Thr Thr Ala Val Gln Thr Pro Thr Ser Gly Glu Pro
 20 25 30

Leu Val Ser Thr Ser Glu Pro Leu Ser Ser Lys Met Tyr Thr Thr Ser
 35 40 45
 Ile Thr Ser Asp Pro Lys Ala Asp Ser Thr Gly Asp Gln Thr Ser Ala
 50 55 60
 Leu Pro Pro Ser Thr Ser Ile Asn Glu Gly Ser Pro Leu Trp Thr Ser
 65 70 75 80
 Ile Gly Ala Ser Thr Gly Ser Pro Leu Pro Glu Pro Thr Thr Tyr Gln
 85 90 95
 Glu Val Ser Ile Lys Met Ser Ser Val Pro Gln Glu Thr Pro His Ala
 100 105 110
 Thr Ser His Pro Ala Val Pro Ile Thr Ala Asn Ser Leu Gly Ser His
 115 120 125
 Thr Val Thr Gly Gly Thr Ile Thr Thr Asn Ser Pro Glu Thr Ser Ser
 130 135 140
 Arg Thr Ser Gly Ala Pro Val Thr Thr Ala Ala Ser Ser Leu Glu Thr
 145 150 155 160
 Ser Arg Gly Thr Ser Gly Pro Pro Leu Thr Met Ala Thr Val Ser Leu
 165 170 175
 Glu Thr Ser Lys Gly Thr Ser Gly Pro Pro Val Thr Met Ala Thr Asp
 180 185 190
 Ser Leu Glu Thr Ser Thr Gly Thr Thr Gly Pro Pro Val Thr Met Thr
 195 200 205
 Thr Gly Ser Leu Glu Pro Ser Ser Gly Ala Ser Gly Pro Gln Val Ser
 210 215 220
 Ser Val Lys Leu Ser Thr Met Met Ser Pro Thr Thr Ser Thr Asn Ala
 225 230 235 240
 Ser Thr Val Pro Phe Arg Asn Pro Asp Glu Asn Ser Arg Gly Met Leu
 245 250 255
 Pro Val Ala Val Leu Val Ala Leu Leu Ala Val Ile Val Leu Val Ala
 260 265 270
 Leu Leu Leu Leu Trp Arg Arg Arg Gln Lys Arg Arg Thr Gly Ala Leu
 275 280 285
 Val Leu Ser Arg Gly Gly Lys Arg Asn Gly Val Val Asp Ala Trp Ala
 290 295 300
 Gly Pro Ala Gln Val Pro Glu Glu Gly Ala Val Thr Val Thr Val Gly
 305 310 315 320
 Gly Ser Gly Gly Asp Lys Gly Ser Gly Phe Pro Asp Gly Glu Gly Ser
 325 330 335
 Ser Arg Arg Pro Thr Leu Thr Thr Phe Phe Gly Arg Arg Lys Ser Arg
 340 345 350
 Gln Gly Ser Leu Ala Met Glu Glu Leu Lys Ser Gly Ser Gly Pro Ser
 355 360 365

Leu Lys Gly Glu Glu Glu Pro Leu Val Ala Ser Glu Asp Gly Ala Val
 370 375 380

Asp Ala Pro Ala Pro Asp Glu Pro Glu Gly Gly Asp Gly Ala Ala Pro
 385 390 395 400

<210> 15

<211> 2745

<212> DNA

<213> Homo sapiens heterogeneous nuclear ribonucleoprotein K

<400> 15

cggcagtcctc gcgcgggtac tgcagcactg ggggtgtcagt tgttggtccg acccagaacg 60
 cttcagttct gctctgcaag gatataataa aactgattgg tgtgcccggt taataaaaaga 120
 atatggaaac tgaacagcca gaagaaacct tccctaacac tgaaccaat ggtgaatttg 180
 gtaaacgccc tgcagaagat atggaagagg aacaagcatt taaaagatct agaaacactg 240
 atgagatggg tgaattacgc attctgcttc agagcaagaa tgctggggca gtgattggaa 300
 aaggaggcaa gaatattaag gctctccgta cagactacaa tgccagtgtt tcagtcccag 360
 acagcagtgg ccccgagcgc atattgagta tcagtgtgta tattgaaaca attggagaaa 420
 ttctgaagaa aatcatccct accttggaag agggcctgca gttgccatca cccactgcaa 480
 ccagccagct cccgctcgaa tctgatgctg tggaatgctt aaattaccaa cactataaag 540
 gaagtgactt tgactgcgag ttgaggctgt tgattcatca gagtctagca ggaggaatta 600
 ttgggggtcaa aggtgctaaa atcaaagaac ttcgagagaa cactcaaacc accatcaagc 660
 ttttccagga atgctgtcct cattccactg acagagttgt tcttattgga ggaaaacccg 720
 atagggttgt agagtgcata aagatcatcc ttgatcttat atctgagtct cccatcaaag 780
 gacgtgcaca gccttatgat cccaattttt acgatgaaac ctatgattat ggtgggtttta 840
 caatgatgtt tgatgaccgt cgcggacgcc cagtgggatt tcccatgctg ggaagagggtg 900
 gttttgacag aatgcctcct ggtcgggggtg ggcgtcccat gcctocatct agaagagatt 960
 atgatgatat gagccctcgt cgaggaccac ctccccctcc tcccgacga ggcggccggg 1020
 gtggttagcag agctcggaat ctctctcttc ctccaccacc accacctaga gggggagacc 1080
 tcatggccta tgacagaaga gggagacctg gagaccgtta cgacggcatg gttggtttca 1140
 gtgctgatga aacttgggac tctgcaatag atacatggag cccatcagaa tggcagatgg 1200
 cttatgaacc acaggggtggc tccgatatg attattccta tgcagggggg cgtggctcat 1260
 atggtgatct tgggtggacct attattacta cacaagtaac tattccc aaa gatttggtg 1320
 gatctattat tggcaaagggt ggtcagcggg ttaaacaat ccgatcatgag tcgggagctt 1380
 cgatcaaaat tgatgagcct ttagaaggat ccgaagatcg gatcattacc attacaggaa 1440
 cacaggacca gatacagaat gcacagtatt tgctgcagaa cagtgtgaag cagtatgcag 1500
 atgttgaagg attctaagtc aagatatttt ttctttttta tagtgtgaag cagtattctg 1560

gaaagttttt ctaagactag tgaagaactg aaggagtcct gcaccttttt ttttttatct 1620
 gcttctgttt aaaaagccaa cattcctctg ctccataggt gtctctgcatt tgagggtgtag 1680
 tgaaatcttt gctgttcacc agatgtaatg ttttagttcc ttacaaacag ggttgggggg 1740
 gggaagggcg tgcaaaaact aacattgaaa ttttgaaaca gcagcagagt gagtggattt 1800
 tatttttctg tattgttggg ggtttaaaaa attcccccca tgtaattatt gtgaacacct 1860
 tgctttgtgg tcaactgtaac atttgggggg tgggacaggg aggaaaagta acaatagtcc 1920
 acatgtccct ggcatctgtt cagagcagtg tgcagaatgt aatgctcttt tgtaagaaac 1980
 gttttatgat ttttaaaata aatttagtga acctattttt ggtggtcatt ttttttttaa 2040
 gacagtcatt ttaaaatggg ggctgaattt cccaaccac ccccaaacta aacactaagt 2100
 ttaattttca gctcctctgt tggacatata agtgcacttc ttgttggaca taggcaaaat 2160
 aacttgccaa acttagttct ggtgatttct tgatggtttg gaagtctatt gctgggaaga 2220
 aattccatca tacatattca tgcttataat aagctgggga ttttttgttt gtttttgcaa 2280
 atgcttgccc ctacttttca acaattttct atgttagttg tgaagaacta aggtggggag 2340
 cagtactaca agttgagtaa tggtagagt atataccaga attctgattg gcagcaagtt 2400
 tattaatcag aataacactt ggttatggaa gtgactaatg ctgaaaaaat tgattatttt 2460
 tattagataa tttctcacct atagacttaa actgtcaatt tgctctagtg tcttattagt 2520
 taaactttgt aaaatatata tatacttgtt tttccattgt atgcaaattg aaagaaaaag 2580
 atgtaccatt tctctgttgt atgttgatt atgttagaat gtttgtgtac aattcaaaaa 2640
 aaaaaagat gaaaaagtt cctgtggatg ttttgtgtag tatcttgga tttgtattga 2700
 tagttaaaat tcacttccaa ataaataaaa caccatgat gctag 2745

<210> 16

<211> 463

<212> PRT

<213> Homo sapiens heterogeneous nuclear ribonucleoprotein complex K

<400> 16

Met Glu Thr Glu Gln Pro Glu Glu Thr Phe Pro Asn Thr Glu Thr Asn
 1 5 10 15

Gly Glu Phe Gly Lys Arg Pro Ala Glu Asp Met Glu Glu Glu Gln Ala
 20 25 30

Phe Lys Arg Ser Arg Asn Thr Asp Glu Met Val Glu Leu Arg Ile Leu
 35 40 45

Leu Gln Ser Lys Asn Ala Gly Ala Val Ile Gly Lys Gly Gly Lys Asn
 50 55 60

Ile Lys Ala Leu Arg Thr Asp Tyr Asn Ala Ser Val Ser Val Pro Asp
 65 70 75 80

Ser Ser Gly Pro Glu Arg Ile Leu Ser Ile Ser Ala Asp Ile Glu Thr
 85 90 95
 Ile Gly Glu Ile Leu Lys Lys Ile Ile Pro Thr Leu Glu Glu Gly Leu
 100 105 110
 Gln Leu Pro Ser Pro Thr Ala Thr Ser Gln Leu Pro Leu Glu Ser Asp
 115 120 125
 Ala Val Glu Cys Leu Asn Tyr Gln His Tyr Lys Gly Ser Asp Phe Asp
 130 135 140
 Cys Glu Leu Arg Leu Leu Ile His Gln Ser Leu Ala Gly Gly Ile Ile
 145 150 155 160
 Gly Val Lys Gly Ala Lys Ile Lys Glu Leu Arg Glu Asn Thr Gln Thr
 165 170 175
 Thr Ile Lys Leu Phe Gln Glu Cys Cys Pro His Ser Thr Asp Arg Val
 180 185 190
 Val Leu Ile Gly Gly Lys Pro Asp Arg Val Val Glu Cys Ile Lys Ile
 195 200 205
 Ile Leu Asp Leu Ile Ser Glu Ser Pro Ile Lys Gly Arg Ala Gln Pro
 210 215 220
 Tyr Asp Pro Asn Phe Tyr Asp Glu Thr Tyr Asp Tyr Gly Gly Phe Thr
 225 230 235 240
 Met Met Phe Asp Asp Arg Arg Gly Arg Pro Val Gly Phe Pro Met Arg
 245 250 255
 Gly Arg Gly Gly Phe Asp Arg Met Pro Pro Gly Arg Gly Gly Arg Pro
 260 265 270
 Met Pro Pro Ser Arg Arg Asp Tyr Asp Asp Met Ser Pro Arg Arg Gly
 275 280 285
 Pro Pro Pro Pro Pro Gly Arg Gly Gly Arg Gly Gly Ser Arg Ala
 290 295 300
 Arg Asn Leu Pro Leu Pro Pro Pro Pro Pro Arg Gly Gly Asp Leu
 305 310 315 320
 Met Ala Tyr Asp Arg Arg Gly Arg Pro Gly Asp Arg Tyr Asp Gly Met
 325 330 335
 Val Gly Phe Ser Ala Asp Glu Thr Trp Asp Ser Ala Ile Asp Thr Trp
 340 345 350
 Ser Pro Ser Glu Trp Gln Met Ala Tyr Glu Pro Gln Gly Gly Ser Gly
 355 360 365
 Tyr Asp Tyr Ser Tyr Ala Gly Gly Arg Gly Ser Tyr Gly Asp Leu Gly
 370 375 380
 Gly Pro Ile Ile Thr Thr Gln Val Thr Ile Pro Lys Asp Leu Ala Gly
 385 390 395 400

Ser Ile Ile Gly Lys Gly Gly Gln Arg Ile Lys Gln Ile Arg His Glu
 405 410 415
 Ser Gly Ala Ser Ile Lys Ile Asp Glu Pro Leu Glu Gly Ser Glu Asp
 420 425 430
 Arg Ile Ile Thr Ile Thr Gly Thr Gln Asp Gln Ile Gln Asn Ala Gln
 435 440 445
 Tyr Leu Leu Gln Asn Ser Val Lys Gln Tyr Ser Gly Lys Phe Phe
 450 455 460

<210> 17
 <211> 1144
 <212> DNA
 <213> Homo sapiens Pur (pur-alpha)

<400> 17
 cgactgaggc ggcgggcgga gcggcaggcg gcggcggcgc ggcagcggag cgcagcatca 60
 tggcggaccg agacagcggc agcgagcagg gtggtgcggc gctgggttcg ggcggctccc 120
 tggggcaccc cggctcgggc tcaggctccg gcggggcgcg tggtgccggc gggggcgggc 180
 gcggcagtgg cggcggcggc ggcgggggccc caggggggct gcagcacgag acgcaggagc 240
 tggcctccaa gcgggtggac atccagaaca agcgcttcta cctggacgtg aagcagaacg 300
 ccaagggccg ctctctgaag atcgccgagg tgggcgcggg cggcaacaag agcgcctta 360
 ctctctccat gtcagtggcc gtggagtcc gcgactacct gggcgacttc atcgagcact 420
 acgcgcagct gggccccagc cagccgcggc acctggccca ggcgcaggac gagccgcgcc 480
 gggcgctcaa aagcgagttc ctggtgcgcg agaaccgcaa gtactacatg gatctcaagg 540
 agaaccagcg cggccgcttc ctgcgcattc gccagacggt caaccggggg cctggcctgg 600
 gctccacgca gggccagacc attgcgctgc ccgcgcaggg gctcatcgag ttccgtgacg 660
 ctctggccaa gctcatcgac gactacggag tggaggagga gccggccgag ctgcccaggg 720
 gcacctcctt gactgtggac aacaagcgct tcttcttoga tgtgggctcc aacaagtacg 780
 gcgtgtttat gcgagtgagc gaggtgaagc ccacctatcg caactccatc accgtgccct 840
 acaaggtgtg ggccaagtgc ggacacacct tctgcaagta ctcgaggag atgaagaaga 900
 ttcaagagaa gcagagggag aagcgggctg cctgtgagca gcttcaccag cagcaacagc 960
 agcagcagga ggagaccgcc gctgccactc tgctactgca gggtgaggaa gaaggggaag 1020
 aagattgatc aaacagaatg aaacccccac acacacacac atgcatacac acacacacac 1080
 agccacacac acagaaaata tactgtaaag aaagagagaa aataaaaagt taaaaagtta 1140
 aaaa 1144

<210> 18
 <211> 322
 <212> PRT

<213> Homo sapiens purine-rich element binding protein A (PURA)

<400> 18

Met	Ala	Asp	Arg	Asp	Ser	Gly	Ser	Glu	Gln	Gly	Gly	Ala	Ala	Leu	Gly	
1				5				10						15		
Ser	Gly	Gly	Ser	Leu	Gly	His	Pro	Gly	Ser	Gly	Ser	Gly	Ser	Gly	Gly	
			20					25					30			
Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Gly	
		35					40					45				
Gly	Ala	Pro	Gly	Gly	Leu	Gln	His	Glu	Thr	Gln	Glu	Leu	Ala	Ser	Lys	
	50					55					60					
Arg	Val	Asp	Ile	Gln	Asn	Lys	Arg	Phe	Tyr	Leu	Asp	Val	Lys	Gln	Asn	
65					70					75				80		
Ala	Lys	Gly	Arg	Phe	Leu	Lys	Ile	Ala	Glu	Val	Gly	Ala	Gly	Gly	Asn	
				85					90					95		
Lys	Ser	Arg	Leu	Thr	Leu	Ser	Met	Ser	Val	Ala	Val	Glu	Phe	Arg	Asp	
			100					105					110			
Tyr	Leu	Gly	Asp	Phe	Ile	Glu	His	Tyr	Ala	Gln	Leu	Gly	Pro	Ser	Gln	
	115					120						125				
Pro	Pro	Asp	Leu	Ala	Gln	Ala	Gln	Asp	Glu	Pro	Arg	Arg	Ala	Leu	Lys	
	130					135					140					
Ser	Glu	Phe	Leu	Val	Arg	Glu	Asn	Arg	Lys	Tyr	Tyr	Met	Asp	Leu	Lys	
145					150					155				160		
Glu	Asn	Gln	Arg	Gly	Arg	Phe	Leu	Arg	Ile	Arg	Gln	Thr	Val	Asn	Arg	
				165				170						175		
Gly	Pro	Gly	Leu	Gly	Ser	Thr	Gln	Gly	Gln	Thr	Ile	Ala	Leu	Pro	Ala	
			180					185					190			
Gln	Gly	Leu	Ile	Glu	Phe	Arg	Asp	Ala	Leu	Ala	Lys	Leu	Ile	Asp	Asp	
	195						200					205				
Tyr	Gly	Val	Glu	Glu	Glu	Pro	Ala	Glu	Leu	Pro	Glu	Gly	Thr	Ser	Leu	
	210					215					220					
Thr	Val	Asp	Asn	Lys	Arg	Phe	Phe	Phe	Asp	Val	Gly	Ser	Asn	Lys	Tyr	
225					230					235				240		
Gly	Val	Phe	Met	Arg	Val	Ser	Glu	Val	Lys	Pro	Thr	Tyr	Arg	Asn	Ser	
				245					250					255		
Ile	Thr	Val	Pro	Tyr	Lys	Val	Trp	Ala	Lys	Phe	Gly	His	Thr	Phe	Cys	
			260					265					270			
Lys	Tyr	Ser	Glu	Glu	Met	Lys	Lys	Ile	Gln	Glu	Lys	Gln	Arg	Glu	Lys	
		275					280					285				
Arg	Ala	Ala	Cys	Glu	Gln	Leu	His	Gln	Gln	Gln	Gln	Gln	Gln	Gln	Glu	
	290					295						300				
Glu	Thr	Ala	Ala	Ala	Thr	Leu	Leu	Leu	Gln	Gly	Glu	Glu	Glu	Gly	Glu	
305					310					315					320	

Glu Asp

<210> 19
<211> 22
<212> DNA
<213> Synthetic oligonucleotide (CD43 PyRo SS)

<400> 19
gggccactt cctttcccct tg 22

<210> 20
<211> 16
<212> DNA
<213> Synthetic oligonucleotide (CD43 PyRo SSUB)

<220>
<221> misc_feature
<222> (9)..(10)
<223> bromouracil

<220>
<221> misc_feature
<222> (13)..(15)
<223> bromouracil

<220>
<221> misc_feature
<222> (20)..(21)
<223> bromouracil

<220>
<221> misc_feature
<222> (23)..(23)
<223> biotin

<400> 20
gggccacccc ccccgb 16

<210> 21
<211> 22
<212> DNA
<213> Synthetic oligonucleotide (CD43 Mut-11)

<400> 21
gggccactt ccttcatata tg 22

<210> 22
<211> 20
<212> DNA
<213> Synthetic oligonucleotide (NS-SS)

<400> 22
gagttagctc actcattagg 20

<210> 23
<211> 21
<212> DNA
<213> Synthetic oligonucleotide (LUC-2)

<400> 23
atagccttat gcagttgctc t 21

<210> 24
<211> 39
<212> DNA
<213> Synthetic oligonucleotide (GeneRacer RNA Oligo)

<220>
<221> misc_feature
<222> (5)..(5)
<223> bromouracil

<220>
<221> misc_feature
<222> (21)..(21)
<223> bromouracil

<220>
<221> misc_feature
<222> (26)..(26)
<223> bromouracil

<220>
<221> misc_feature
<222> (31)..(31)
<223> bromouracil

<220>
<221> misc_feature
<222> (39)..(39)
<223> bromouracil

<400> 24
cgacggagca cgaggacacg acaggacgaa ggagagaaa 39

<210> 25
<211> 54
<212> DNA
<213> Synthetic oligonucleotide (GeneRacer Oligo dT Primer)

<400> 25
gctgtcaacg atacgctacg taacggcatg acagtgtttt tttttttttt tttt 54

<210> 26
<211> 23
<212> DNA
<213> Synthetic oligonucleotide (GeneRacer 5' Primer)

<400> 26
cgactggagc acaggacac tga 23

<210> 27
<211> 27
<212> DNA
<213> Synthetic oligonucleotide (GeneRacer 5' Nested Primer)

<400> 27
ggacactgac catggactga aggagta 27

<210> 28
<211> 33
<212> DNA
<213> Synthetic oligonucleotide (LUC-4)

<400> 28
cactacggta ggctgcgaaa tggttcatact gtt 33

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)